

Where does plant-based meat fit in the UPF conversation?

A detailed guide to key topics and proof points for health professionals and stakeholders in protein diversification.



About this resource

This document was co-created by the Good Food Institute Europe (GFI Europe) and the Physicians Association for Nutrition (PAN), two non-profit organisations working to support the transition towards healthier, more sustainable diets.

It is designed to offer an objective and detailed but also accessible overview of the current discourse to stakeholders in protein diversification, including healthcare professionals, policymakers, researchers, health and sustainability NGOs and patient organisations.

Authors:

- **Amy Williams** (Nutrition Lead, GFI Europe)
- **Dr Johanneke Tummers** (PAN International Research Fellow, School for Moral Ambition)
- **Dr Roberta Alessandrini** (Director, Dietary Guidelines Initiative, PAN International)

Conflict of interest statement: None of the authors or their affiliated organisations have any financial relationships with food companies or producers of plant-based meat.

This document is not intended to offer individual health guidance.

Using this resource


This resource is intended to offer an evidence-based overview of research and key considerations relating to plant-based meat, ultra-processed foods (UPFs) and nutrition in the European context.

The sections divide the current debate into key topics, and explore the context and evidence for each.

To improve accessibility for non-specialist audiences, certain technical words (found in **bold**) are defined in the glossary on page 49.

Contents:

- 4. [Introduction: Using this resource](#)
- 6. [How is plant-based meat made and how is it produced?](#)
- 12. [What are ultra-processed foods?](#)
- 21. [Existing research on the effects of swapping conventional meat for plant-based meat](#)
- 26. [Research available on UPFs and its relevance to plant-based meat](#)
- 35. [Summary, conclusions and recommendations](#)
- 40. [Appendix](#)
- 49. [Glossary](#)
- 54. [References](#)



What is the ultra-processed foods (UPFs) debate?

Image: Vegfather

Research on ultra-processed foods (UPFs) shows that people with diets high in UPFs tend to have lower **diet quality** and higher rates of various diseases compared to people whose diets contain the least ([Lane et al. 2024](#)).

These **epidemiological** studies highlight that people with an increased disease risk are eating more convenient, cheap, high-calorie, and low-nutrient foods like sugary drinks, processed meats, cakes, and biscuits, while consuming fewer nutrient-rich foods like fruits, vegetables, legumes, and whole grains.

Most UPFs are high in calories and low in nutrients, but the term is not defined by nutritional criteria, and some UPFs have good nutritional value.

Research is ongoing to explore whether those eating the most UPF have higher disease risk because the food they eat tends to be higher in salt, fat and sugar or if some of this is caused by processing itself ([Dicken et al NCT05627570](#), Hall et al NCT 05290064).

Because we don't know how much of the effect is caused by processing and how much is nutrient profile, there is debate about how UPF classifications should be used in policy and nutrition education, especially when assessing individual foods rather than overall diets.



How does this relate to plant-based meat?

Image: Quorn

Plant-based meat is often considered UPF, but differs from the broader group in several ways.

Unlike most UPFs, plant-based meat is generally low in sugar and saturated fat, high in protein and a source of fibre ([Espinosa et al 2024](#)).

Observational (real-world) **epidemiological studies** that look at how different kinds of UPF affect health find that while people who eat the most UPF overall have increased health risks, risk is not uniform: some UPF types like processed meat and sugary drinks have particularly elevated risk, while others are not associated with increased risk ([Dicken et al 2024](#), [Cordova et al 2023](#)).

Experimental trials suggest replacing conventional meat with plant-based meat can reduce **LDL (bad) cholesterol** and bodyweight ([Fernández-Rodríguez et al 2025](#)) – which are opposite to findings in the UPF group as a whole.

Plant-based meat could support shifts towards healthier, more sustainable dietary patterns ([Messina et al 2023](#)). However, because of the prevalent UPF discourse, their health profile is often misunderstood by the public. Health professionals, policymakers and public health communicators should collaborate to address this.

What is plant-based meat, and how is it produced?

In this section:

- Plant-based meat is designed to replicate the taste and texture of animal meat using plant-based ingredients.
- The nutritional composition and processing level of plant-based meat varies by product and by country.
- The main points in production requiring processing are making the protein base, **texturisation** and added ingredients.
- Each of these can be done using higher or lower degrees of processing.
- Unlike most UPFs, plant-based meat is usually high in protein, a source of fibre (unlike conventional meat), low in saturated fat and low in sugar ([Espinosa et al 2024](#)).
- Inconsistent **fortification** with key nutrients and further reductions in salt content are two areas where the nutritional value of plant-based meat could be improved.

Topic: What is plant-based meat?

Statement	Further detail
Plant-based meat is designed to replicate the taste and texture of animal meat.	- Plant-based meat aims to replicate the taste and texture of conventional animal meat, differentiating it from traditional high-protein plant-based foods like seitan, tofu, and Textured Vegetable Protein (TVP). - Plant-based meat is made from plants (or sometimes fungi). The level of processing used varies by product (<i>see page 8</i>).
Many different plant-based meat products are available in Europe today, with varied nutritional profiles and techniques used to make them.	- A pork hot dog has a very different nutritional profile from a chicken breast, and this variation is also true in plant-based meat. - Different processing approaches can have advantages and disadvantages. For instance, processing can enhance digestibility of protein and micronutrients but it can also increase salt content (Cargo-Froom et al 2023, Samtia et al 2020, Xia et al 2024). - Soy, wheat and pea protein concentrates and isolates are some of the most common bases for plant-based meat, made by extracting the protein from raw plant ingredients. This is called fractionation and can be done in several ways. - Fermentation is also used to make protein ingredients, generally this involves less processing. Mycoprotein (protein from fungi like mushrooms and yeast) is already a common ingredient in the UK, and newer fermentation techniques can create meatier textures and umami flavours. Texturisation is also needed to create a meaty texture. Extrusion is a texturisation process commonly used to make plant-based meat, as well as many other widely eaten foods like shaped pasta and cereals.
Plant-based meat has particular public health potential to help reduce widespread overconsumption of processed meat, without requiring major behavioural change.	- Plant based meat is sometimes viewed as a niche food for those already following plant-rich diets like vegetarians, but its primary potential for public health lies in broader mainstream adoption. - It has the largest potential for the many people who enjoy meaty meals, eat less fibre and more processed meat than recommended, and don't want to completely overhaul their diets, preferring options that fit within their existing daily routines. - Increased availability of tasty, affordable, nutritious plant-based meat designed to appeal to these people could help improve diet quality, and make plant-based foods more accessible and less intimidating.

Overview: common processes used for high-protein plant-based foods

Protein base

A core ingredient with high **protein density** is made, either by extracting protein from plants (eg soy, peas), or by growing fungi or other microorganisms with high protein content.

Separating the protein out from the raw ingredients is sometimes called **fractionation**, and the process used determines how much of the **food matrix** from the base ingredients remains in the end product.

Texturisation

This protein base is then mixed with other ingredients (see page 9) and undergoes one of several possible processing steps to recreate the texture of conventional meat.

It is not easy to tell based on information on the packaging what **texturisation** processes have been used to make foods.

Lower



Higher

Processing
level

Examples:

- Biomass fermentation
- Washing wheat flour
- Soy curds
- **Concentrates and isolates**
 - Dry **fractioning** (separation by weight)
 - Wet **fractioning** (separation by solubility)

For more details on these processes, see appendix slide 41.

Examples:

- Solid-state fermentation
- Pressing/moulding
- 3D printing
- **Extrusion**
- Fibre spinning

For more details on these processes, see appendix slide 42.

Overview: other ingredients used in plant-based meat

UPFs on average have longer ingredient lists, so this metric is a common shorthand to identify them. However, ingredient lists are generally poorly understood by non-specialist audiences.

For example, the public are often wary of ‘**emulsifiers**’, even though this term describes the ingredient’s function rather than its composition, which varies. For instance, minimally processed ingredients like egg yolk are common **emulsifiers**. Other **emulsifiers** like **lectins** are present (but not added and therefore not labelled) in whole plant foods and animal meat ([Pryme et al 2021](#), [Radhakrishnan et al 2022](#)), but can cause concern when listed as an ingredient.

All food ingredients approved by Europe’s rigorous food safety processes have publicly available safety assessments on the [EFSA website](#). Approvals require **experimental trials** and long-term health monitoring.

Structural ingredients

Used to help replicate the texture of conventional meat.

A wide variety are used with different nutrient profiles and processing levels – ranging from plant fibres to ingredients like **methylcellulose**, lectins and gums, which are used as **emulsifiers** and binders to help mix the ingredients together without separating.

Flavours and colours

These range from familiar ingredients like salt, herbs and beetroot extract to **artificial ingredients** made using chemical processes.

Traditional fermentation can be used to create complex umami flavours with low processing, and **precision fermentation** ingredients like **haem**, already used outside Europe, can deliver authentic meaty flavours and **bioavailable** iron.

Fats and oils

A wide variety of fats and oils are used in different plant-based meat products, with varying processing levels and health attributes.

These range from unsaturated fats like rapeseed and olive oil, to saturated fats like coconut, palm, and **hydrogenated oils**, or a mix of these.

Fortification

Fortification with nutrients like B12 and iodine is common in animal feed, leading to increased and consistent presence of certain nutrients in animal products.

Conversely, **fortification** is inconsistent in plant-based meat, meaning content of key nutrients like vitamin B12 varies. This should be improved.

Topic: The nutritional composition of plant-based meat

Point

Significant variation exists between products and categories, but in general plant-based meat has a positive nutritional profile.

Further detail

Compared to animal-based counterparts* and EU/UK **health claim** definitions, plant based meat is on average:

- Similar or slightly lower **caloric density**.
- A source of fibre, while conventional meat is not.
- High in protein. Similar to conventional meat in terms of % of calories from protein, but slightly lower per 100g.
- Neither low nor high in fat, similar to conventional meat.
- Low in saturated fat, unlike conventional meat.
- Low in sugar, like conventional meat.
- In most categories neither low nor high in salt.
- It should be noted that while the average seen here is roughly similar, large variation exists within categories: plant-based meat is higher in salt than unseasoned, unprocessed conventional meat, but similar or lower compared to processed conventional meat.
- See Appendix (A3-A8) for full breakdowns by product type and nutrient




Figure created using median data from studies of nutritional composition of plant-based and conventional meat. Calories, fibre, saturated fat, sugar and salt taken from [Espinosa et al 2024](#). Protein was not included in Espinosa et al, so protein figures were calculated based on values from other recent papers in Germany, Spain, Italy and the UK ([Grea et al 2023](#), [Heras-Delgado et al 2023](#), [Cutroneo et al 2022](#), [Alessandrini et al 2021](#)).

*Unless otherwise specified, conventional meat comprises both processed and unprocessed products grouped together. Full breakdowns by product type can be found in the appendix.

Topic: The nutritional composition of plant-based meat

Point	Further detail
Plant based meat is sometimes, but not always, fortified with key nutrients.	• Animal feed is commonly fortified with key nutrients like B vitamins, calcium, iodine, long chain omega 3s and vitamin D (Neill et al 2021, ADHB 2025, Heutgen 2010). This fortification supports animal growth, but also boosts concentrations and consistency in animal products. Consequently, in European diets today fortified animal feed is a major source of several key nutrients. It is therefore important that plant-based options contain equivalent fortification. • Fortification is used in certain staple foods like flour and margarine in several countries, and in some it is even mandatory (European Commission 2006, FAO 2021). • In plant-based meat however, fortification is inconsistent both on the country and product level, which should be improved to increase the potential for these products to meet similar nutritional needs. • Fortification rates are improving over time, and government guidelines such as those in the Netherlands (Voedingcentrum 2025) can significantly accelerate this: ○ A study in 2023 found 55% of Dutch meat alternative products were fortified (van Haperen 2023), which had risen to 75% in a similar study in 2024 (Gallani et al 2024). ○ Increases have been seen elsewhere but most countries have far lower prevalence of fortification overall. For instance, fortification rates in the UK grew from 5% to over 25% between 2021 (Alessandrini et al 2021) and 2024 (Gallani et al 2024). • While fortification via animal feed is not listed as an ingredient, it is mandatory to specify it when added directly to food, thereby lengthening ingredients lists. • Public understanding of UPF has made many people wary of longer ingredient lists, potentially penalising more nutritionally complete plant-based meat options.

What are Ultra-processed foods?


In this section:

- UPFs are most commonly classified using the Nova framework ([Monteiro *et al* 2019](#)), which groups foods into four categories:
 - Nova 1 – foods made with minimal processing
 - Nova 2 – ingredients used in home cooking
 - Nova 3 – somewhat processed foods
 - Nova 4 – heavily (or ‘ultra-’) processed foods.
- The Nova framework does not consider nutritional composition, it separates foods based on how they are made and used.
- Plant-based meat is often considered Nova 4, but is very different from the most widely eaten UPFs both in terms of its nutritional profile and the dietary patterns it tends to fall into.
- The Nova framework can be interpreted in two ways, and plant-based meat fits differently in each.
 - **Epidemiological**, identifying changing dietary patterns that are leading to increased rates of diet-related ill health. This is well researched.
 - **Food-level**, to assess healthfulness of individual foods. This research is still preliminary, and not yet strong enough to warrant interventions independent of well understood factors like nutrient profile ([Lancet Gastroenterology & Hepatology 2024](#)).

Topic: Definitions of food processing using the Nova framework (Monteiro *et al* 2019)

	Nova 1: Unprocessed or minimally processed foods	Nova 2: Processed culinary ingredients	Nova 3: Processed foods	Nova 4: Ultra-processed foods
Definition	- Minimal processing includes the removal of inedible or unwanted parts of a food source and very basic preservation like drying. - Nothing should be added to the original food in this category at the point of sale.	- This category was broadly created to bridge the gap between ready-made and home cooked foods, comprising the ingredients like salt, herbs, sugar, vinegar and oil that are typically added during cooking but do not constitute the bulk of the meal. - These foods are rarely eaten by themselves and unlike the other three categories are not grouped by the way in which they are made, but rather by the way in which they are used.	- Foods from group one that have been processed and/or combined with foods from group two, or group one or two foods that have been further processed into a final product that is ready to cook or eat. - There are generally fewer processing steps and ingredients in processed foods, although there can be overlap in the processes and ingredients used, and grey areas in dividing foods between the two.	- Foods with many ingredients in them and multiple processing steps that could not be recreated in a conventional kitchen. - These foods are often (though not necessarily) high in salt, sugar and fat, low in fibre and calorie dense. - These foods are generally convenient and quick to prepare, cheap, tasty, and often have plastic packaging.
Examples	Fruits, vegetables, beans, mushrooms, tea, eggs, plain cuts of fresh meat (including red meat, white meat and seafood).	Salt, peper, herbs, cooking oil, sugar, vanilla extract, bicarbonate of soda.	Sourdough bread, canned chickpeas, tofu, tempeh, bacon, salted crisps, roasted nuts, yoghurt, cheese, pasta, mayonnaise.	Energy drinks, hot dogs, chocolate bars, shaped salty snacks such as puffs or hoops, biscuits, breakfast cereals, milkshakes, protein powder, pre-packaged bread, plant-based meat.

Overview: Plant-based meat does not neatly fit with many of the characteristics often associated with UPFs – particularly compared to the conventional processed meat it often replaces.



Comparison based on GFI Europe analyses of median data from studies into macronutrient profiles of processed meat and plant-based meat in Spain ([Heras-Delgado et al 2023](#)), the Netherlands ([van Haperen 2023](#)), Sweden ([Bryngelsson et al 2022](#)), the UK ([Alessandrini et al 2021](#)) and Germany ([Grea et al 2023](#)). Subjective categories of 'convenient' and 'could not be made in a conventional kitchen' were determined based on standard Nova definitions. Does not include unprocessed conventional meat.

Topic: Food processing and the origins of the Nova framework

Statement	Further detail
The origins of the Nova framework	• The most frequently used definition of UPFs is based on the Nova framework, developed by Brazilian epidemiologist Carlos Monteiro in 2009. • He saw growing rates of obesity and diet-related ill health, and linked them to the rapidly changing eating habits in Brazil where, like many other places in the world, traditional home cooked foods were eaten less and less, replaced by cheap, mass produced convenience foods. He also saw that large food companies making these mass-produced foods were gaining more control over the food system. • Monteiro argued that these foods, which were designed to be tasty, low-cost and were heavily marketed, were making people less healthy while increasing profits for food companies. • This created a vicious cycle, where fresh, high-quality foods became less accessible, and people lost basic cooking skills needed to prepare healthy meals with vegetables, legumes, and other whole foods. • Monteiro linked these trends and suggested that food processing itself – which makes food cheaper, tastier, and easier to consume – was a key reason for declining diet quality.
What specifically is meant by the term 'processing'?	• Based on this epidemiological observation at the population level, Monteiro sought to define 'processing' on the food level (<i>full definitions detailed on slide 13</i>). • This is a challenge, because processing covers a huge range of varied techniques, is a fundamental part of most food preparation, and is also demonstrably not linear in its impact on the 'healthiness' of a food: both over and undercooking food can undermine its nutritional value, and some cooking techniques like steaming can enhance it. • Certain processing techniques like cooking, pasteurising, and fermenting have been used for centuries to improve the taste, safety and nutritional value of foods. Likewise, many unprocessed foods can be harmful in excess, like pork belly, which is high in saturated fat, or brazil nuts which have high levels of selenium. • Consequently, the definitions of the four Nova categories are descriptive rather than quantitative, and have evolved over time, broadly seeking to draw a 'line of best fit'.

Topic: The two interpretations of the Nova framework

Statement	Further detail
<u>Interpretation 1: Epidemiological.</u> What we eat (and by extension our health) is shaped by social influences and our food environment. Dietary patterns high in UPF are associated with poor health outcomes. Exploring these dietary patterns may offer insight into socio-political levers that can help improve people's diet quality.	• Most UPF research to date uses this epidemiological interpretation of Nova. • Nova's inventor Professor Monteiro specialises in the epidemiology of obesity, and the Nova framework was initially developed on this basis to try and explain rising obesity rates. • UPF studies using this framework typically use long-term, real world datasets. They generally compare the outcomes of those people whose diets contain the most UPFs, with those whose diets contain the least. • Plant-based meat makes up a miniscule proportion of food eaten (0.2%). Meanwhile, the foods contributing the largest proportion of UPF calories in peoples diets come from foods already understood to be unhealthy in excess like pastries, buns, cakes and biscuits (10.8%) (Rauber et al 2024). Given these studies look at diets as a whole rather than individual foods, it is much more likely that high consumption of these less healthy foods – not the very small amount of plant-based meat eaten – drove the outcomes observed.
<u>Interpretation 2: Food-level profiling.</u> Processing level may alter the healthfulness of a food independent of nutrient profile. Understanding the impacts of different processes can help improve the nutritional value of foods.	• At the time of writing (May 2025), only a limited amount of research exists on the extent to which processing causes poor health outcomes independent of nutritional factors like salt, fat and sugar content, nor which processes are most harmful. Several studies exploring this are underway (Dicken et al NCT05627570, Hall et al NCT05290064) which assess the impacts of diets high in UPF versus diets high in minimally processed foods, using UPFs that meet national health recommendations / do not contain high levels of nutrients of concern. • RCTs to date have suggested that harms of high UPF diets do at least in part come from nutritional factors such as caloric density, coupled with more processing-related considerations such as food texture and 'hyper-palatability' encouraging over eating. (Hall et al 2019, Hamano et al 2024). • We already know that the nutritional profiles of different plant-based meat products vary considerably (as with most types of food). The findings of this research on the ways in which processing can either reduce or improve the nutritional quality of food will likely be valuable to further enhance plant-based meat, eg, ways of better preserving some of the beneficial compounds in whole plant foods, optimising the types of fibre and fat present, or further reducing caloric density.

Topic: the Nova framework from a socio-political perspective

Statement	Further detail
The epidemiological view of Nova also points to several social and economic trends at play in our food system, with relevance for plant-based meat.	• A problem highlighted by proponents of the Nova framework is the decline of small independent food producers and businesses in an increasingly globalised food system. ◦ This has already been seen in the conventional meat sector, with smaller family farms struggling to compete with larger companies in a globalised market. ◦ Europe currently has a diverse and innovative ecosystem of plant-based meat companies, with many small startups and medium-sized independent players, alongside brands owned by larger food companies. ◦ Amid the turbulence of the prevailing economic climate, however, smaller companies are particularly exposed to supply chain shocks and cost of living increases. • It is therefore essential that public funding to support open-access research into these foods is increased, support is broadened for affordable access to scaling infrastructure, and barriers to entry are reduced for smaller, independent players to ensure the current diversity continues. • This funding should also support exploration of plant-based meat's potential to support public goods, such as: ◦ Further ways to enhance its nutritional profile. ◦ Procurement pathways that provide robust demand for diverse crops grown by local producers to support rural communities and enhance food security. ◦ Support for the proliferation of affordable products that are accessible to the groups of people most likely to benefit from them, rather than expensive niche products for small groups of people.

Topic: Strengths and limitations of the Nova framework


Statement	Further detail
Strengths of the Nova framework	• Nova characterises the prevailing trends in our food system driving less healthy dietary patterns, and their relationship with demographic and social factors. • The Nova framework clearly resonates with many people, and has effectively rejuvenated interest in well-worn but still very important nutritional guidance to increase consumption of vegetables, legumes and whole grains, and reduce dependence on convenient snack foods that are high in calories and low in nutritional value. It also encourages home cooked foods, which are generally (though not always) a healthier choice. (Mills et al 2017). • Many proponents of the Nova framework point to its holistic nature as an intuitive pathway towards a healthier lifestyle more broadly, which is a fundamental part of successfully achieving lasting improvements in diet.
Limitations of the Nova framework	• Particularly among untrained public audiences, there is widespread confusion as to the definition of UPFs and the real-world implications of research for individuals. Behavioural changes prompted by warnings against UPFs have not been widely tested, and it is unclear whether Nova consistently incentivises the most beneficial dietary shifts. • It is true that the average Nova 4 food is less healthy than the average Nova 1 food, but when accounting for nutrient composition, the differences in healthfulness between Nova categories is not well evidenced, and specific information about individual foods are more informative than membership in these broader categories. • The few available datasets that have been used to try and validate Nova on a food-by-food basis are generally not designed to sort food by processing level. They also often rely on food diaries, which are generally an unreliable data source (Bajunaid et al 2025), that were taken before the Nova system was invented, and lack the granular details necessary to group foods consistently. • Nova does not adequately distinguish between harmful, harmless, and helpful processing that food can undergo, which is particularly confusing for non-specialist audiences. • Nova does not adequately account for nutritional variation derived from processing steps that take place in home cooking, such as boiling versus steaming vegetables (Lee et al 2017), or hotter versus cooler frying temperatures (Mavlanov et al 2025).

Topic: Alternatives to the Nova framework

Statement	Further detail
Other frameworks have been developed to characterise UPFs.	Several other metrics such as SIGA (Christodoulou et al 2020) and IUFoST (Ahrne et al 2025) have been developed aiming to offer more discrete (clear-cut) definitions for ultra-processing and tie them to nutritional characteristics, but these are at a far earlier stage of research (UK SACN, 2023).
Nutritionally-based frameworks that have been subject to robust scientific validation are also used to evaluate healthfulness of foods.	- Many tools to help people evaluate food healthiness are used across Europe in food labels or government scanning apps such as the Nutriscore, NutrInform Battery, Nordic Keyhole, Schijf van Fijv and UK Traffic Light systems. - These have been developed to categorise foods on the basis of nutrient content, broadly recommending lower intakes of nutrients of concern – fat, salt, sugar – and increased intake of desirable nutrients like fibre. - The factors chosen as the basis of these systems are similar, based on well established national nutritional recommendations. There is however variation between each in how much each nutrient contributes to the overall score (eg fat versus sugar). Nutriscore is the most widely researched and so used here, but each has strengths and weaknesses. - The lowest scoring foods in these systems frequently overlap with the UPF category, with one study of French products finding 63% of UPF fell in the ‘poor’ D-E categories, while only 13% fell in the ‘good’ A-B (Sarda et al 2024, see page 20). However, there are areas of divergence as UPFs with good nutrient profiles like packaged wholemeal bread and low-sugar fortified cereals are distinguished from well-established ‘unhealthy foods’. - Plant-based meat generally performs well against these criteria. One study surveying the nutriscore and Nova categories of products available in Spain found that 68% of plant-based meat products fell within the ‘good’ A-B range, while only 26% of conventional meat counterpart products did. (Heras-Delgado et al 2023, see page 20). - Results of studies exploring health outcomes linked to diets high in low-scoring foods according to Nutri-score find similar results to UPF studies, such as elevated cardiovascular disease risk (Deschasaux-Tanguy et al 2024). - This highlights the need for experimental data to understand the balance between nutrient profiles and processing level in their contributions to health outcomes.

Independent of processing level, most (but not all) foods in the UPF category have a poor nutritional profile. Plant-based meat is a significant outlier, and generally performs favourably compared to both the processed conventional meat it replaces and the UPF group overall.

Comparison of Nutri-score ratings of plant-based meat compared to their conventional meat counterparts and the UPF group as a whole*



Reproduction of data published in [Heras-Delgado S. et al \(2023\)](#) and [Sarda B et al \(2024\)](#).

*Note: No single European study was available comparing Nova categorisations for plant-based meat, its conventional counterparts, and the UPF group as a whole, so overall UPF group data was taken from a separate study of products available in France. As such, the overall UPF group nutriscore findings may not be directly comparable, but are consistent with other research on Nutriscore and Nova alignment and so likely indicative.

Existing research* on the effect of swapping conventional meat for plant-based meat

In this section:

- A small but growing body of evidence highlights several opportunities for public health through greater support for replacing conventional (particularly processed) meat with plant-based meat ([Espinosa et al 2024](#)).
- In particular, a **systematic review** and **meta analysis** of randomised controlled trials found a significant drop in **LDL (bad) cholesterol** and reductions in weight when conventional meat was swapped for plant-based meat ([Fernández-Rodríguez et al 2025](#)).
- Other studies have found **diet quality**, **microbiome** and gut health benefits ([Farsi et al 2023](#), [Bottin et al 2016](#)).
- These results make sense given plant-based meat has higher fibre and lower saturated fat than conventional meat ([Hartley et al 2016](#), [Hooper et al 2020](#)).
- These findings are very different from those seen in studies of health outcomes linked to high intake of foods in the overall UPF category.
- More research is still needed.

Existing research on the effect of swapping conventional meat for plant-based meat

Point	Further detail
Plant-based meat can help expand uptake of plant-based foods alongside other strategies.	• Initiatives supporting greater intake of whole plant foods are sorely needed. However, these initiatives alone may not be enough to meet environmental and health targets within timeframes needed. • Many people want to eat more plants, but not always for the same reasons – some are motivated by health, others by environment, others by a mix of these and other factors (Reuzé et al 2023). These people also come from a range of circumstances and backgrounds, meaning it's highly likely that different approaches will work better for different people, and driving change will require a combination of complementary approaches. • Despite being a much more recent invention, in the UK plant-based meat volume sales are over 7 times that of tofu, and 5 times that of traditional bean burgers (GFI Europe, 2024), suggesting it may appeal to broader groups of people. • Plant-based meat and similar foods like plant-based dairy, in addition to offering fortification opportunities, require only small behavioural shifts for adoption. They therefore make it easier for people to initiate and maintain more plant-centric dietary patterns (Alae-Carew, 2022).
Studies exploring the impacts of replacing conventional meat with plant-based meat are generally in line with what we would expect based on their nutritional profiles.	• Several experimental trials have explored the impact of replacing conventional meat with plant-based meat. Findings are broadly positive, particularly for reduction of LDL (bad) cholesterol and weight loss (Fernández-Rodríguez et al 2025, see page 24 and 25). • Three independent systematic reviews, of which one was a meta analysis, were published in 2024 synthesising the findings from these studies, from researchers based in Spain, the UK and Canada (Fernández-Rodríguez et al 2025, Espinosa et al 2024, Nagra et al 2024). • Most studies were small, but results were generally consistent, and match what would be expected based on their fibre content and fat profile compared to conventional meat (Hartley et al 2016, Hooper et al 2020). • While more, longer, and larger trials would be helpful to better understand these opportunities, there is already compelling evidence that plant-based meat can support the shift towards healthier, more sustainable diets.

Existing research on the effect of swapping conventional meat for plant-based meat

Point


Clinical trials have shown promising findings in several areas.

Further detail

Trials suggest plant-based meat may help:

- Reduction of LDL (bad) cholesterol (*see page 24*).
- Weight loss in people with overweight (*see page 25*).
- Benefits for gut health and lower colorectal cancer risk ([Farsi et al 2023](#)).
- Maintain **muscle protein synthesis (MPS)** rates in older people similarly to conventional meat ([Domić et al 2024](#)). This contrasts with findings of lower MPS rates from meals with matching calorie and protein content using only whole food plant-protein sources compared to meat ([Pinckaers et al 2024](#)). This difference is likely due to improved **protein quality** achieved through certain processing techniques.

Key findings from interventional trials exploring the health impacts of replacing animal meat with plant-based meat



Further research is needed to better understand the generalisability of results to date.

Key areas for future research should include:

- Comparisons of impact by protein ingredient base, and the contribution of key factors such as fibre or **protein density** to outcomes. Most current studies use **mycoprotein**-based and soy-based products, with the strongest benefits seen in **mycoprotein**.
- More studies into the relative **bioavailability** of protein and key **micronutrients** in plant-based versus conventional meat.
- Studies to understand the effectiveness of plant-based meat in supporting adherence to healthier dietary shifts compared to other approaches, and whether benefits persist on longer time frames.
- Studies in more diverse populations to understand if variation exists in impacts between different demographic groups.
- Explorations in populations with specific dietary needs such as athletes, older people, or those looking to maintain lean mass during weight loss, for whom nutrient density, protein and fibre are key considerations.

Overview: Fernandez-Rodriguez et al, 2024. The first major systematic review and meta analysis of randomised controlled trials on plant-based meat.



Source: Rubén Fernández-Rodríguez, Bruno Bizzozero-Peroni, Valentina Díaz-Goñi, Miriam Garrido-Miguel, Gabriele Bertotti, Alberto Roldán-Ruiz, Miguel López-Moreno. Plant-based meat alternatives and cardiometabolic health: a systematic review and meta-analysis. The American Journal of Clinical Nutrition 2025. Volume 121, Issue 2, February 2025, Pages 274–283 doi: <https://doi.org/10.1016/j.ajcnut.2024.12.002>

Number of participants: Seven randomised controlled trials reported across eight research papers, covering 369 adults.

Study design: Data was pooled from included trials, exploring changes in indicators of cardiovascular health associated with the substitution of conventional meat with plant-based meat for 1-8 weeks, to understand consistency and extent of outcomes across trials.

Outcomes examined: Total, LDL (bad) and HDL (good) cholesterol, body weight, fasting blood glucose, and blood triglycerides.

Key findings: Swapping conventional meat for plant-based meat was associated with **significant reductions in LDL-cholesterol, total cholesterol and body weight**. The overall reduction in LDL cholesterol* was 12% (-0.25 mmol/L), and largest in **mycoprotein** interventions (-37 mmol/L).

No significant differences were observed in other outcomes studied.

*For context, reductions of 0.2-0.4 mmol/L represent ‘moderate’ reductions. This places plant-based meat on the higher end of effectiveness as a dietary intervention to lower LDL ([Schoeneck et al 2021](#)). The European Society of Cardiology recommend LDL should be ‘as low as possible’ for all populations, and are particularly important for those with high risk of **cardiovascular disease**. Reductions achievable through diet are smaller than with medicines, which can reduce levels by over 60% ([ESC/EAS, 2019](#)). If the reductions seen in these trials were maintained over a long period of time they could contribute to reduced **cardiovascular disease** risk ([Penson et al, 2020](#)).

Overview: RE-MAP trial, 2022. A large, publicly funded RCT on the effectiveness of plant-based meat as an intervention to reduce overconsumption of meat.



Number of participants: 115 British adults who regularly ate meat.

Study design: Participants were randomised to receiving free plant-based meat and guidance on the environmental and health advantages of eating more plant-based foods for a period of 4 weeks, and after another 4 weeks were checked on to review behaviour change following the intervention. The control received no intervention or advice on dietary change.

Outcomes examined: The primary outcome was reduction in meat consumption per day, with secondary outcomes exploring whether changes persisted after the trial finished, and health **biomarkers** such as weight, cardiovascular risk factors and diet nutrient composition.

Key findings: Those in the plant-based meat arm ate an average of 63g less meat per day during the intervention and also experienced statistically significant weight loss of half a kilo. This brought consumption by those in the intervention group down from an excess of meat eaten at baseline (130g/day) to within the recommended daily amount (70g/day or less). At follow-up 4 weeks after the intervention finished, participants were still eating 39g less per day compared to baseline. No other significant changes were observed. No changes were seen in the control group at either time point.

Source: Bianchi F, Stewart C, Astbury NM, Cook B, Aveyard P, Jebb SA. Replacing meat with alternative plant-based products (RE-MAP): a randomized controlled trial of a multicomponent behavioral intervention to reduce meat consumption. *Am J Clin Nutr*. 2022 May 1;115(5):1357–1366. doi: [10.1093/ajcn/nqab414](https://doi.org/10.1093/ajcn/nqab414). PMID: 34958364; PMCID: PMC9071457.

Research available to date* on UPF and its relevance to plant-based meat

In this section:

- Several features of datasets used in UPF research limit their relevance to plant-based meat:
 - Most rely on food diaries taken before Nova and most modern plant-based meat existed, and lack the detail needed to categorise newer foods like plant-based meat.
 - Plant-based meat makes up a tiny proportion of UPF eaten, and other UPF foods probably drove observed outcomes.
- **Experimental trials** ([Hall et al 2019](#), [Hamano et al 2024](#)) suggest high **caloric density** and low fibre content (which don't apply to plant-based meat) are at least partially behind the health impacts of high-UPF diets.
- Publicly funded bodies in France, Germany, Spain, and the UK ([ANSES 2025](#), [DGE 2023](#), [Aesan 2020](#), [SACN 2025](#)) have independently found insufficient evidence to support policies targeting UPFs that have a good nutritional profile.
- There are large discrepancies between the findings of research to date and the high volume of poor quality media reporting on the topic.

*As of 20 May 2025

Topic: Research to date on UPFs and its relevance to plant-based meat

Statement	Further detail
Most of the findings to date come from observational (or ‘real-world’) studies, and focus on patterns at the population level.	• It can be a challenge to tease out the long-term impacts of specific foods and different dietary patterns because the kinds of diseases associated with poor diet tend to take years to develop. • Real-world evidence, which tracks data from people going about their daily lives over long periods of time, is therefore often used to research dietary patterns, and comprises most UPF studies to date. • These studies show groups of people who eat the most UPF, and the least minimally processed food, are more likely to develop a number of negative health outcomes, from heart disease to cancer to depression. • However, this data can only tell us about dietary patterns, not individual foods, limiting relevance to plant-based meat: ◦ Because most UPFs are high in calories, salt, fat and sugar, and low in nutrients and fibre, these studies are at risk of confounding, and cannot tell us how much of the health impacts are caused by processing, and how much by these nutritional features we already know negatively impact health. ◦ Plant-based meat, which generally has a good nutritional profile, makes up only 0.2% of calories eaten in datasets like UK Biobank (Rauber et al 2024), meaning findings on UPFs overall are unlikely to apply. ◦ Population-level data is less able to explore specific questions in smaller, non-standard groups. Plant-based meat does not contribute a significant proportion of calories in average high UPF diets, and UK Biobank data suggests those who eat the most plant-based meat generally follow non-standard dietary patterns such as pescetarian or vegetarian (suppl material. Chang et al 2024). Those following these diets in the UK Biobank dataset generally had lower risk of cancer (Parra-Soto et al 2022) and cardiovascular disease (Petermann-Rocha et al 2021), unlike those eating the most UPF generally. ◦ To prove what is causing the observed outcomes and separate out these factors, studies using other designs such as randomized-controlled trials (RCTs) would be needed. • This is a challenge because RCTs are more expensive and take longer to conduct, while the harms of poor diet are pressing, and people are very interested in any research they think can help them improve their health.


Topic: Most commonly used datasets for UPF studies

Statement	Further detail
What datasets are used in real-world UPF studies?	• Many of the landmark UPF studies to date are based on datasets from large health mapping projects set up to monitor trends in key health metrics over long periods of time. Three major European examples of these datasets include: ○ The EPIC cancer database. ■ A database following over half a million enrollees in 10 European countries, started in the 1990s. ■ At enrollment, each person completed in-depth health questionnaires, including on diet, lifestyle and other factors like health status, medications and basic measurements. ■ Follow-up was then conducted at varying points to measure changes over time. ○ The Navarra SUN study. ■ A Spanish database started in 1999 to explore the nutritional benefits of the Mediterranean diet, which now has over 20,000 enrollees, mostly alumni from the University of Navarra. ■ Anyone can register, after which they complete a baseline questionnaire covering food frequency, health status, medicine usage and lifestyle. ■ A shorter follow-up questionnaire is then taken every two years to track changes over time. ○ The UK Biobank ■ This is the largest biomedical database in the world, founded in 2006, with over half a million UK adults enrolled. ■ Anonymised information on genomes and electronic health records are collected alongside the baseline questionnaires. ■ Additional smaller groups of participants have also contributed other information like body, brain and heart scans, activity data from wearables, and questionnaires on behaviour, diet, mental health and pain management. • Other commonly used datasets from outside Europe include the US NHANES database and Nurses Health Study.

Topic: Available research into UPF and health

Statement	Further detail
What are the challenges of using these datasets to explore UPF in relation to plant-based meat?	• Certain limitations of these datasets make it particularly difficult to use them to assess plant-based meat: ○ The food intake data predates Nova, and so lacks granular detail essential for categorisation. ■ In food frequency questionnaires used by these databases, ‘meat alternatives’, if recorded at all, were not grouped by processing level making it impossible to separate traditional products like tofu (Nova 3) from plant-based meat (Nova 4) (Riboli et al 2003 (EPIC), Oxford WebQ (Biobank)). ■ Few people know much about how plant-based meat is made – including UPF researchers. Plain mycoprotein-based products made using biomass fermentation and freezing would generally fall in Nova 3, and yet they are typically assumed to be made with extrusion and placed in Nova 4. ○ Food intake data is usually from before many modern plant-based meat options existed. ■ Studies are looking for changes over long time periods, so most UPF studies use the baseline food frequency questionnaires to map UPF intake. ■ This means in studies using the EPIC cohort, UPF intake data is at least 25 years old (Dicken et al 2024), and in those using the UK Biobank it is at least 10 years old (Chang et al 2023). While some plant-based meat products do predate this, the large expansion of plant-based meat as a category began around 2019, long after any of these data were collected. ○ The people in these studies get a vanishingly small proportion of their calories from plant-based meat, making it impossible to separate the impact of plant-based meat from more commonly eaten UPFs. ■ In the Biobank dataset, just 0.2% of calories came from ‘plant-based alternatives to meat’, which also included tofu (suppl material. Rauber et al 2024). By comparison, 16% were from sweet snacks and drinks like pastries, cakes, biscuits, soft drinks and confectionery, and 9% were from UPF animal sourced foods like milkshakes, processed meat and dairy-based puddings (Rauber et al 2024). ■ Several of these larger UPF groups are independently associated with negative health outcomes (Nguyen et al 2022, Feng et al 2021).

The average increased risk of developing cardiovascular disease, coronary heart disease or stroke for those with diets high in UPF as a whole (light blue), and those with diets high in UPF excluding processed meat and sugary drinks (dark blue)




“The diverse nutritional composition within these products warrants the need to deconstruct the ultra-processed food classification for a nuanced understanding of their impact on cardiovascular health. **Our findings suggest that soft drinks and processed meats should be particularly discouraged**, given their consistent adverse association with cardiovascular disease, coronary heart disease, and stroke.”

Reproduction using data from: [Mendoza, Kenny et al. Ultra-processed foods and cardiovascular disease: analysis of three large US prospective cohorts and a systematic review and meta-analysis of prospective cohort studies, The Lancet Regional Health – Americas, 2024, Volume 37, 100859](#)

Topic: Available research into UPF and health

Statement	Further detail
Real world studies have found that different kinds of UPF have different associated health risks, with sugary drinks and processed meat being the two subgroups most strongly associated with harm.	- Studies breaking down ultra-processed foods by sub-category have found that processed meat, savoury snacks and sugary drinks drove a significant proportion of the adverse health associations with UPF, with processing levels in other subgroups seeming to have far weaker, if any, negative links (Cordova et al 2024, Dicken et al 2024). - A systematic review of several trials, which normalised UPF intake to remove the influence of foods already linked with increased risk on the category as a whole, found that with processed meat and sugary drinks removed, the risk associations with cardiovascular disease and stroke disappeared (Mendoza et al 2024, see page 30). Another looking at diabetes found that processing level had far weaker risk associations than specific food groups like processed meat and sugary drinks taken individually (Mendoza et al 2025). - This suggests that while the epidemiological lens of viewing Nova may hold, more nuance is likely needed when using it to profile individual foods – particularly those with generally favourable nutritional profiles like plant-based meat.
Two interventional trials have been completed, and a partial readout from a third has been released to date (March 2025). All three suggest that nutritional characteristics of these foods do factor into the health impacts of UPF.	- While there are many real-world studies, many are based on the same people in the same 5-6 datasets (see page 28) and have very similar methodologies and research questions. This makes it very important to see the results from new studies using a wider variety of study designs to gain a deeper understanding – particularly RCTs. - Several randomised controlled trials are underway seeking to test whether the associations between UPF consumption and negative health outcomes are still seen when normalising for factors such as nutrient composition, and to date two have been published (Hall et al 2019, Hamano et al 2024 see pages 32 and 33). - These preliminary data have teased out more detail on what specifically about UPF may drive the patterns seen in real world studies, highlighting certain specific aspects of processing that may be driving increased health risks: - The higher caloric density of diets high in UPF (ie number of calories per 100g). - The hyper-palatability and softer texture of many UPF foods, encouraging excess consumption. - The lower fibre content of high UPF diets, and the type and format of fibre (ie in food or as a drink supplement). - These trials are still small and only cover very short time periods, but they suggest UPF's health impacts are at least in part due to nutritional factors like caloric density and fibre content – both of which plant-based meat performs well on.

Summary: Hall et al, 2019 – the first randomised controlled trial on UPF



Number of participants: 20

Study design: 20 healthy American participants were randomised to either UPF or non-UPF based diets for 14 days. Once complete, there was a break and the two groups swapped for another 14 days, so both tried both the UPF and non-UPF menu. The meals themselves were matched for calories and macronutrients (although distribution throughout the meal varied), but all were far larger than a single portion, so participants would eat until they were full.


Was plant-based meat included? No

Outcomes examined: The primary outcome explored was differences in daily calorie intake, with change in body weight from baseline, change in body fat mass, perceived food pleasantness, eating speed and satiety as secondary outcomes.

Key findings: “Energy intake was greater during the ultra-processed diet, with increased consumption of carbohydrate and fat, but not protein. **Weight changes were highly correlated with energy intake**, with participants gaining 0.9 ± 0.3 kg during the ultra-processed diet and losing 0.9 ± 0.3 kg during the unprocessed diet.” No differences were found in participant ratings of food pleasantness, eating speed, liver fat, or fullness. Body fat measures were inconclusive, with gains in both fat mass and fat-free mass in the UPF diet and losses of both in the non-UPF diet.

Summary: Hamano et al, 2024 – the second published RCT on UPF

33



Number of participants: 9

Study design: Nine overweight Japanese male participants were randomised to either UPF or non-UPF based diets for seven days. Once complete, there was a break and the two groups swapped for another seven days, so both tried both the UPF and non-UPF menu. The meals themselves were matched for calories and macronutrients (although distribution throughout the meal varied), but all were far larger than a single portion, so participants would eat until they were full.

Was plant-based meat included? No

Outcomes examined: The primary outcome explored was changes in body weight, with daily average energy intake and chewing frequency as secondary outcomes.

Key findings: Unlike in the Hall study, weight gain was observed in both the UPF and non-UPF arms, however during the UPF period, **participants gained 1.1 kg more weight and consumed 813.5 kcal more per day compared with during the non-UPF period.** Regarding the chewing frequency, the number of chews per calorie was significantly lower during the UPF period.

Topic: Available research into UPF and health

Statement	Further detail
Several national health bodies have found that there is no sufficient evidence to warrant policy interventions on the basis of Nova category alone, recommending that, if used, it is combined with other complementary factors that account for nutritional profile.	- Many European governments agree on the need to address the growing problems linked to poor diets. However, independent, publicly funded investigations from multiple European countries all agree that the evidence is not strong enough to recommend policy or public health education based on processing level alone: - France's ANSES report, published in January 2025. - Germany's DGE systematic review, published December 2023. - The UK's SACN report, published in July 2023 and updated in April 2025, and the UK House of Lords report on Food, Diet and Obesity, published October 2024. - Spain's Aesan report, published in 2020. The 2023 Nordic Nutrition recommendations also do not recommend use of UPF over existing frameworks. - However, the general public remains confused by the seeming disconnect between these conclusions and the high volume of news stories on the topic, and what the findings mean in practice. - Better communication on this topic is sorely needed to help people make better food choices.
The research to date is unlikely to tell us much about plant-based meat.	- It is clear that the narrative surrounding UPF has given many people a misleading impression of the nutritional profile of plant-based meat, even though there is good evidence it could be a helpful tool to support healthier and more sustainable choices. - Foods like processed meat, which plant-based meat is designed to replace, appear to play a large role in driving increased health risks associated with UPFs. - Making processed foods healthier is likely a key part of effectively improving diet-related ill health on a population level. In parallel, another key component lies in making healthy whole foods like vegetables, legumes and beans more accessible and affordable. These two strategies likely complement one another.

Summary, conclusions and recommendations

Summary

Plant-based meat makes up a vanishingly small proportion of food eaten in both epidemiological studies and RCTs on UPFs, and also has a very different nutritional profile to most UPFs. It is therefore unlikely this research can tell us much about plant-based meat's health impacts.

- In the UK Biobank dataset, plant-based meat made up just 0.2% of calories eaten ([Rauber et al 2024](#)).
- Most UPFs are high in fat, sugar, salt and **caloric density**, and low in fibre. **Caloric density** and low fibre content have been identified as the most likely mechanisms for health impacts seen in existing RCTs looking at UPFs. On average, plant-based meat is not high in fat, sugar or salt, is a source of fibre, and does not have high **caloric density**. This suggests studies on the UPF group as a whole likely offer little insight into plant-based meat, and will likely be misleading as to its health impacts.
- RCTs exploring the effects of plant-based meat as a replacement for conventional meat have generally found positive effects, which are in line with what we would expect based on their nutritional profile, in particular their fibre content and fat profile. These RCTs are better quality evidence.

There is room for improvement in the nutritional profile of plant-based meat, particularly more consistent fortification and reduction of nutrients of concern such as salt.

- A lot of variation currently exists in the nutritional profile of plant-based meat available in Europe. In particular, rates of **fortification** are currently inconsistent, although they are improving over time. Salt content is currently moderate, and most people could benefit from reducing salt intake.
- The best available evidence nevertheless suggests that replacing processed meat with plant-based versions could have medically relevant health benefits, most notably reductions in LDL cholesterol.

People are currently eating more than recommended daily intakes of meat, and more options to help reduce this has several public health advantages.

- There is good evidence linking over-consumption of processed meat with higher risk of colorectal cancer, the second leading cause of cancer death in Europe. The negative impact of processed meat is also highlighted by UPF studies.
- Publicly funded research to make tasty, affordable plant-based meat options that can out-compete the conventional products driving this over consumption may help support adherence to healthier, more sustainable diets.
- This would have broader public health benefits, such as reducing the risks of emergent zoonotic diseases and antimicrobial resistance, both of which are exacerbated by modern intensive farming practices ([Kelbrick et al 2023](#), [Hayek 2022](#)).

Conclusions

Research on ultra-processed foods has broadened the **epidemiological** understanding of diet-related ill health, and unlocked political will to drive much-needed change in our food system.

Plant-based meat is an accessible option with particular opportunities as a replacement for processed meat – which has been identified by research into UPF as one of the subcategories most strongly associated with increased health risks.

Although trials have identified several key features of UPFs as a whole that are likely to play a large role in the negative outcomes observed (high **calorie density**, low fibre and **hyper-palatability**); plant-based meat has a very different nutritional profile from most UPFs, and these metrics do not generally apply to plant-based meat.

With current meat consumption in Europe above recommendations for both public and planetary health, a diverse range of strategies are likely required to support the necessary increase in consumption of plant-based foods. There is no single path to achieving this, and multiple complementary strategies are likely needed. Support for tasty, affordable plant-based meat and encouraging greater uptake of whole plant foods are two such strategies, and each likely has differing appeal to different demographic groups.

On this basis, proponents of the Nova framework and of protein diversification should be incentivised to identify where each can have the most impact building a healthier, more sustainable food system, and collaborate to target their work accordingly.

Recommendations

Public health and nutrition professionals should challenge misconceptions on processing and plant-based meat.

- UPF research is poorly understood by the general public, and misconceptions are common.
- This is compounded by limited familiarity with plant-based meat and its nutritional profile, with many assuming it is high in salt, sugar and fat like the majority of UPF.
- Better communication on these points to improve accuracy and reduce sensationalism is sorely needed.
- Greater emphasis should be placed on the fact that there are multiple pathways to healthier lifestyles – and the most effective interventions are those that can be adhered to.

Researchers should focus on diversifying the evidence base for UPF and plant-based meat.

- Interventional studies (eg randomised controlled trials) are needed to understand what features of UPF drive harm, and how to effectively address the impacts of the most harmful foods.
- Such studies are also needed to explore which qualities of plant-based meat are behind observed beneficial outcomes to help guide further improvements.
- Behavioural research is needed to understand the best levers to achieve widespread dietary improvement, and how to effectively target the groups most affected by diet-related ill health.
- Food diaries and food frequency questionnaires used in nutritional **epidemiology** should be designed to capture plant-based meat consumption.

National bodies should create guidelines to help people select healthier options, and increase consistency across products.

- This includes guidelines on features like **fortification**, fibre and salt, where variation across plant-based meat products exists.
- Products meeting these positive thresholds should also be included in national dietary guidance, as is the case in the Netherlands.
- Guidelines that support more accessible pathways to healthy dietary patterns should also be explored.


Learn more about plant-based meat and health



Appendix

A1: common processes used for high-protein plant-based foods

Protein base



Biomass fermentation. Used for plant-based meat. Certain kinds of fungi that grow small fibres – called filamentous fungi – are grown in a fermenter. They are then siphoned off and warmed before being separated from the water, giving a minimally processed ingredient high in **soluble fibre** and **complete protein** ([Quorn, 2017](#)).

Wheat flours. Used for **seitan**. Wheat flour is made into a dough then washed, leaving the protein behind. While the protein concentration is high, it is not complete, and much of the fibre is removed.

Soy curds. Used for tofu. Dried soybeans are soaked, blended and cooked. The resulting soy milk is separated from the fibre pulp. A **coagulant** is then added to curdle the milk. This works by changing the pH to match its **isoelectric point**, causing the protein to **precipitate** into curds (like in cheesemaking). These solid curds are then removed and pressed. These have complete protein and while some fibre is lost, more is kept than in **seitan**.

Concentrates and isolates. Used for plant-based meat. Several methods can be used to separate protein from other constituents in raw plant ingredients (**fractionation**), resulting in different nutritional characteristics. **Concentrates** retain more of the original **food matrix**, including fiber and beneficial plant compounds like polyphenols, while **isolates** are nearly pure protein. Two main methods are used:


- **Dry fractionation** (used to make **concentrates**) separates protein by particle size and weight. First, the raw ingredient is dried and milled. During milling, protein-rich particles tend to be smaller than those containing more starch or fiber. Air is then used to separate these smaller, protein-dense particles from the larger starch- or fiber-rich ones, forming the concentrate. This method is energy-efficient and produces fewer byproducts, but is usually lower in protein and harder to texturise.
- **Wet fractionation** (used to make **concentrates** and **isolates**) separates protein based on solubility. The raw ingredient is ground into a meal and, for oil-rich sources like soybeans and chickpeas, defatted. If already a liquid, the pH can be changed to precipitate the protein curd out of solution. If not, the protein is first dissolved using alkaline agents. The resulting protein solution is then extracted, and precipitated into a curd as in the other processes. This curd is then washed, filtered or centrifuged, and dried to produce a powdered **isolate** with over 90% protein content.

Mixing with
other
ingredients

See slide 9 for more
information on other
commonly used
ingredients.

A2: common processes used for high-protein plant-based foods


Texturisation



A3: calorie density in plant-based versus conventional meat

Point	Further detail
Plant-based meat tends to have lower calorie density .	- Choosing foods with lower calories by weight can reduce overall calorie intake (Robinson, et al 2022). - Randomised controlled trials exploring mechanisms for UPF impacts on health suggest their higher average calorie density is one of the primary drivers of observed weight gain (Hall et al 2019, Hamano et al 2024). - Plant-based meat does not have a high calorie density, in contrast to many other UPFs. - On average, plant-based meat products have similar or fewer calories per 100g than conventional meat.

Average calories per 100g in plant-based and conventional meat



Reproduced from GFI Europe report, [Plant-based meat and health \(2023\)](#). Synthesis of median data from studies of nutritional composition of plant-based and conventional meat in Germany ([Grea et al 2023](#)), the Netherlands ([van Haperen 2023](#)), Spain ([Heras-Delgado et al 2023](#)), Sweden ([Bryngelsson et al 2022](#)) and the UK ([Alessandrini et al 2021](#)).

A4: Fibre in plant-based versus conventional meat


Point

Plant-based meat tends to be a source of fibre

Further detail

- There is growing evidence that high fibre intake helps maintain a healthy weight ([Thompson et al 2017](#)) and reduces **biomarkers** associated with **cardiovascular disease** ([Hartley et al 2016](#)), echoed in studies of real world data ([Threapleton et al 2013](#), [Stephen et al 2017](#)).
- Initial studies have also suggested fibre intake may influence gut and **microbiome** health ([So et al 2018](#)).
- Most Europeans do not eat enough fibre in their diet.
- Plant-based meat therefore offers an opportunity to increase fibre content in parts of meals that previously had none, with only minor tweaks to the recipe.
- Fibre content is another area where plant-based meat differs from average UPFs.
- Plant-based meat is a source of fibre. Conventional meat is not.

Average fibre per 100g in plant-based and conventional meat



Reproduced from GFI Europe report, [Plant-based meat and health \(2023\)](#). Synthesis of median data from studies of nutritional composition of plant-based and conventional meat in Germany ([Grea et al 2023](#)), the Netherlands ([van Haperen 2023](#)), Spain ([Heras-Delgado et al 2023](#)), Sweden ([Bryngelsson et al 2022](#)) and the UK ([Alessandrini et al 2021](#)).

A5: protein density in plant-based versus conventional meat


Point

Plant-based meat tends to be high in protein

Further detail

- Plant-based meat comfortably meets the EU 'high protein' definition, with a percentage of calories from protein similar to that of conventional meat in most categories except lean fillets.
- The processing used to make plant-based meat can improve **protein density**, and **quality** relative to their raw ingredients ([Manzanilla-Valdez et al 2024](#)). More research to quantify specific impacts on these factors by processing method is needed.
- Protein density** and **quality** are particular considerations for helping those with lower appetites (eg, older people), aiming for high protein intake within tight calorie thresholds (eg, athletes), or otherwise in **calorie deficit** (eg, trying to lose weight) maintain muscle mass ([Domić et al 2025](#), [Jäger et al 2017](#), [Janssen et al 2023](#)).

Average % of calories from protein in plant-based and conventional meat



Synthesis of median data from studies of nutritional composition of plant-based and conventional meat in Germany ([Grea et al 2023](#)), the Netherlands ([van Haperen 2023](#)), Spain ([Heras-Delgado et al 2023](#)), Sweden ([Bryngelsson et al 2022](#)) and the UK ([Alessandrini et al 2021](#)).

A6: saturated fat in plant-based versus conventional meat


Point

Plant-based meat tends to be low in saturated fat

Further detail

- There is good evidence that long-term reduction of saturated fat intake can reduce risk of cardiovascular events like heart attacks or strokes ([Hooper et al 2020](#)).
- Plant-based meat usually contains low levels of saturated fat, and significantly less than animal-based counterparts.

Average saturated fat per 100g in plant-based and conventional meat



Reproduced from GFI Europe report, [Plant-based meat and health \(2023\)](#). Synthesis of median data from studies of nutritional composition of plant-based and conventional meat in Germany ([Grea et al 2023](#)), the Netherlands ([van Haperen 2023](#)), Spain ([Heras-Delgado et al 2023](#)), Sweden ([Bryngelsson et al 2022](#)) and the UK ([Alessandrini et al 2021](#)).

A7: sugar in plant-based versus conventional meat


Point

Plant-based meat tends to be low in sugar

Further detail

- There is good evidence that high sugar consumption, particularly from sugar-sweetened drinks ([Santos et al 2022](#)), is associated with increased risk of **cardiovascular disease** and other chronic diseases ([Huang et al 2023](#)).
- Plant-based meat and conventional meat are both low in sugar.
- While both are 'low sugar' foods, in absolute terms plant-based meat contains slightly more.
- This is another key area where plant-based meat is significantly different from many of the most frequently consumed UPF food groups such as sugary drinks, cakes, biscuits and milkshakes.

Average sugar per 100g in plant-based and conventional meat



Reproduced from GFI Europe report, [Plant-based meat and health \(2023\)](#). Synthesis of median data from studies of nutritional composition of plant-based and conventional meat in Germany ([Grea et al 2023](#)), the Netherlands ([van Haperen 2023](#)), Spain ([Heras-Delgado et al 2023](#)), Sweden ([Bryngelsson et al 2022](#)) and the UK ([Alessandrini et al 2021](#)).

A8: salt in plant-based versus conventional meat


Point

Plant-based meat tends to have moderate salt content

Further detail

- Most Europeans exceed the guideline maximum daily salt intake of 5-6g (2-2.4g sodium) ([European Commission 2021](#), [Kwong et al 2022](#)).
- Excess salt intake is discouraged due to links with **high blood pressure**, which increases **cardiovascular disease** risk ([Graudal et al 2020](#)).
- Plant-based meat's salt content varies by country and product.
- Plant-based meat usually has moderate salt levels: more than unseasoned fresh meat but similar or less than processed meat ([Espinosa et al 2024](#)).
- A 2023 German study found plant-based salami had half the salt of conventional salami, but burgers were similarly salty ([Grea et al 2023](#)).
- RCTs replacing conventional meat with plant-based meat do not find any impact on blood pressure ([Fernández-Rodríguez et al 2025](#)).

Average salt per 100g in plant-based and conventional meat



Synthesis of median data from studies of nutritional composition of plant-based and conventional meat in Germany ([Grea et al 2023](#)), the Netherlands ([van Haperen 2023](#)), Spain ([Heras-Delgado et al 2023](#)), Sweden ([Bryngelsson et al 2022](#)) and the UK ([Alessandrini et al 2021](#)).

Glossary

Artificial ingredients

The EU does not have a legal definition for this term, but it is generally understood to mean ingredients that are synthesized, as opposed to being derived from a plant or animal.

Bioavailability

The proportion of a given nutrient in a food that the body can absorb for use in everyday functions.

Biomarkers

A biological molecule found in blood, other body fluids, or tissues that is a sign of a normal or abnormal process, or of a condition or disease.

Calorie deficit

A person is considered in calorie deficit when they use more energy than they gain from food, leading to weight loss as the body uses up energy stored as fat or muscle tissue to compensate.

Calorie density

Also known as energy density. Defined as the number of calories per 100g. The European Union does not have a formal definition of high calorie density, however 400 calories per 100g is a frequently used threshold for high calorie density ([Nesta, 2024](#)).

Cardiovascular disease

Medical conditions affecting the heart or blood vessels.

Coagulant

A compound or agent which is added to a liquid to thicken it.

Concentrates

Protein concentrates are ingredients used in plant-based meat that are typically made by milling the raw ingredients and separating out the most protein rich flour grains on the basis of their weight. Compared to isolates, they generally have a lower protein content, but retain more of the original food matrix, including fiber and beneficial plant compounds like polyphenols

Confounding factors

A confounding factor in a study is one that influences both the dependant variable and the independent variable, which if not accounted for can give the impression of an incorrect causal relationship between the two things being looked at. To give an example, a study might compare rates of ice cream consumption and incidence of sunburn in a given population, and see that people are more likely to get sunburned on days that they eat ice cream - ie that there is an association between these two variables. However, common sense tells us that eating ice cream does not cause sunburn. This is because of a confounding factor - sunny weather - which increases the rate of both ice cream consumption and sunburn. If we normalise the dataset to remove the influence of sunny weather (which we know independently causes sunburn) on the results, we would expect the association between eating icecream and sunburn to disappear.

Diet quality

Diet quality is a measure of how well someone's diet aligns with dietary guidelines. High quality diets are balanced and contain the right amount of energy, macro and micronutrients to support good health. There are several tools used to measure diet quality, such as the Healthy Eating Index (HEI).

EFSA

European Food Safety Authority - the EU's food safety regulatory body.

Emulsifiers

Emulsification is a general term referring to processes that allow a stable mixture of two or more liquids that would usually separate (eg oil and water). Emulsifiers facilitate this process. Ingredients with emulsifying properties have useful applications in food. Mayonaise is a widely used example of an emulsion, using egg yolk as an emulsifier to combine oil and vinegar.

Epidemiology

The study of how often diseases occur in different groups of people and why. Epidemiological studies typically take large population-level datasets and look for patterns in those with higher rates of certain diseases to try and gain insight into the drivers of a given disease.

Experimental trials

Experimental trials, also known as interventional trials, are studies where researchers introduce a change in the study population to see how it affects an outcome. The most robust interventional study design is a randomised controlled trial (RCT).

Food matrix

The food matrix is the term used to describe the structure, composition and nutrient balance of a whole food. These structures can result in different biological responses compared to processed versions of the same food. For example, sugars in fruit are absorbed by the body at different rates depending on whether the fruit is being eaten whole or has first been blended into juice, or whether the fruit is raw or cooked.

Fortification

Fortification is when specific nutrients (often those identified as a risk of shortfall in the general population) are added to foods to help make it easier for people to get all the nutrients they need from their diet. Fortification of certain staples like bread are legally mandated in several European countries to support public health.

Fractionation / fractioning

Fractionation is a general term referring to any process that can split out the component parts of a mixture on the basis of their physical properties. For instance, using a centrifuge to separate by weight, or heating to separate by boiling point).

Haem

An iron-containing molecule present in blood cells which is used by the body to transport oxygen around the body. If eaten directly, haem iron is more easily used by the body than non-haem iron.

Health claims

Health claims are regulated terminology that can be used to describe beneficial nutritional properties of given foods, provided they meet certain thresholds. These can be positive thresholds for beneficial nutrients (eg high fibre) or upper limits for nutrients harmful in excess (eg low sugar).

High blood pressure

High blood pressure (also known as hypertension) is a condition where the force of a person's blood as it pumped around the body is consistently high for an extended period of time. Over time, this can cause damage to the blood vessels (particularly small ones like those in the kidneys and eyes), and is linked to various health problems.

Hyper-palatability

This is not a standardised term, but generally refers to the relative ratios of fat, sugar, and salt in a given food item. Foods meeting a certain threshold of two or more of these macronutrients by weight are considered hyper-palatable. It has been suggested that such combinations make foods 'extra delicious' and consequently easier to over-eat.

Hydrogenated oils

Hydrogenated oils are saturated fats that have been fully converted from unsaturated fats using hydrogen, making them solid at room temperature. While saturated fats should not be eaten in excess, they are not to be confused with partially-hydrogenated fats, or artificial 'trans-fats,' which used to be used in some foods like baked goods and margarine. Trans fats are also naturally found in small amounts in red and processed meat. Artificial trans fats are strictly regulated and effectively or completely banned altogether in many countries.

Isoelectric point

The pH at which a molecule, especially a protein or amino acid, has no net electrical charge. The isoelectric point of a protein affects its solubility, and changing the pH to match this point causes it to come out of solution.

Isolates

Protein isolates are ingredients with a very high protein content, typically made using wet fractionation, which splits out the component molecules in plants based on their different solubility. They typically have higher protein content than concentrates, but less of the original food matrix.

LDL (bad) cholesterol

LDL cholesterol, or low-density lipoprotein cholesterol, is sometimes called bad cholesterol because it can build up inside a person's arteries, causing them to harden and narrow (a process called atherosclerosis), limiting blood flow. This can ultimately lead to a heart attack or stroke.

Lectins

Lectins are proteins that bind to carbohydrates, they perform a number of biological roles in the cells of living organisms. There are many different types of lectins, some of which are beneficial for health, while others can be harmful. Some lectins found in particularly high concentrations in legumes and grains have anti-nutritional properties, meaning these foods must undergo processing such as heating or fermenting to make them edible. Some lectins can also limit the body's ability to absorb certain micronutrients like iron and zinc, but again processing can be used to reduce or neutralise this effect. Research is still ongoing to fully understand the various ways in which different kinds of lectin can have positive and negative interactions with various functions within the body.

Limiting amino acids

The essential amino acid found in the shortest supply relative to the amounts needed for protein synthesis in the body. Four amino acids are most likely to be limiting: lysine, methionine, threonine and tryptophan.

Macronutrients

Macronutrients are used by the body for energy and provide many of the building blocks for normal bodily functions. They therefore make up the bulk of all the food we eat. There are three kinds of macronutrient: carbohydrates, fats and proteins.

Methylcellulose

Methylcellulose is an ingredient made from cellulose. It was invented in the 1950s and became widely used as a thickener and emulsifier in various foods beginning in the 1960s. Like cellulose, it is not digestible, non-toxic, and not an [allergen](#).

Microbiome

A microbiome is the community of microorganisms that can usually be found living together in any given habitat. In health terms, this usually refers to the various species of microorganisms that live in our digestive tract.

Micronutrients

Micronutrients are vitamins and minerals that are needed by the body for healthy functioning, but only in very small amounts.

Muscle Protein Synthesis (MPS)

Muscle Protein Synthesis rate describes the rate at which the body builds muscle tissue. It is generally lower in older people.

Mycelium

Mycelium is the root-like structure of a fungus, made up of a network of thin threads, which form the main body of the organism.

Mycoprotein

Mycoprotein is a protein ingredient made from fungi. This can either be microscopic fungi or larger fungi like the protein-rich 'roots' of mushrooms known as mycelium.

Quorn is the most widely available example of mycoprotein available today, and has been used as an ingredient since 1985.

Nordic Keyhole

The Nordic Keyhole is a front-of-pack nutrition label used in Nordic countries that helps consumers choose healthier food options within product groups, based on the Nordic Nutrition Recommendations.

NutrInform Battery

The NutrInform Battery is a front-of-pack label used in Italy (also available in a scanning app for phones), displaying the content of energy, fats, saturated fats, sugars, and salt per serving, along with its contribution to daily dietary requirements, using a "battery" symbol to represent percentages.

Nutriscore

Nutri-Score is a colour-coded, front-of-pack nutritional label designed to make a food product's nutritional value easier to understand, using a scale from A (healthiest) to E (least healthy) to help people make informed choices.

Observational (real-world) studies

A type of research that uses data from people going about their daily lives, in this context looking for emerging patterns in people grouped by the proportion of their diet coming from UPF. These studies often have the advantage of having a large number of participants, but they are not randomised, meaning it can be hard to know the impact of one single factor relative to other inter-related characteristics linked to poor health such as smoking status. As such, they are mainly used to either identify a theory to be tested in an experimental trial, or to confirm if an outcome seen in an experimental trial still persists in imperfect real-world conditions.

Precipitated

A chemistry term describing the process whereby a solid (precipitate) comes out of a liquid solution to form a sediment. It is the opposite process to dissolution, when a solid is dissolved into a liquid to form a solution.

Precision fermentation

Precision fermentation is a process that has been used for decades in food production to produce common ingredients such as rennet – used in cheesemaking. It is a process that leverages the natural abilities of certain microorganisms like yeast to over-produce a desired compound, which can then be isolated and used as needed.

Protein density

Defined as the amount of protein per 100g.

Protein quality

This refers to a combination of both amino acid profile and digestability in proteins. It is typically defined using the PDCAAS, which stands for Protein Digestibility-Corrected Amino Acid Score or the DIAAS which stands for the Digestible Indispensable Amino Acid Score.

Schijf van Fijv

The Schijf van Fijv or 'Wheel of five' is the dietary guidance model adopted by the Netherlands. It separates foods into 5 separate groups, each of which has a defined minimum content threshold for desirable nutrients like vitamins and fibre, and maximum thresholds for nutrients of concern like salt and sugar. Foods meeting these thresholds for its category are included in dietary guidance, which people can check using the 'Kies ik Gezond' mobile phone app.

Seitan

Seitan is a food made from gluten, the main protein of wheat, that has been used as a meat alternative for many centuries. It was first developed hundreds of years ago by Buddhist monks in China and Japan.

Solid-state fermentation

Solid state fermentation grows a target (such as koji used in Tempeh or mycelium from mushrooms) on a solid feedstock without free flowing water.

Soluble fibre

Soluble fiber is a type of dietary fiber that dissolves in water, forming a gel-like substance in the gut which helps to regulate blood sugar and cholesterol levels.

Textured vegetable protein (TVP)

Textured vegetable protein (TVP) is a dried protein ingredient that was invented in the 1960's, often coming in the form of small 'mince' pieces or chunks.

Texturisation

Processes that are used to create a desired texture in a foodstuff.

UK Traffic Light system

The UK traffic light food label system uses red, amber, and green colors to indicate whether a food is high, medium, or low in fat, saturated fat, sugars, and salt, designed to help people understand food labels and make healthier choices.

Umami

Umami is the 'savory' flavour in foods. It is frequently associated with foods like stock, meat and fermented foods.

References

Page 4

Lane M M, Gamage E, Du S, Ashtree D N, McGuinness A J, Gauci S et al. Ultra-processed food exposure and adverse health outcomes: umbrella review of epidemiological meta-analyses BMJ 2024; 384 :e077310 doi:10.1136/bmj-2023-077310

Dicken S, Makaronidis J, van Tulleken C, Jassil FC, Hall K, Brown AC, Gandini Wheeler-Kingshott CAM, Fisher A, Batterham R. UPDATE trial: investigating the effects of ultra-processed versus minimally processed diets following UK dietary guidance on health outcomes: a protocol for an 8-week community-based cross-over randomised controlled trial in people with overweight or obesity, followed by a 6-month behavioural intervention. BMJ Open. 2024 Mar 11;14(3):e079027. doi: 10.1136/bmjopen-2023-079027. PMID: 38471681; PMCID: PMC10936475.

Page 5

Sarah Nájera Espinosa, Genevieve Hadida, Anne Jelmar Sietsma, Carmelia Alae-Carew, Grace Turner, Rosemary Green, Silvia Pastorino, Roberto Picetti, Pauline Scheelbeek, Mapping the evidence of novel plant-based foods: a systematic review of nutritional, health, and environmental impacts in high-income countries, Nutrition Reviews, 2024;, nuae031, doi: <https://doi.org/10.1093/nutrit/nuae031>

Samuel J. Dicken, Christina C. Dahm, Daniel B. Ibsen, Anja Olsen, Anne Tjønneland, Mariem Louati-Hajjig, Claire Cadeaug, Chloé Marquesg, Matthias B. Schulzeh, Franziska Jannaschh, Ivan Baldassarik, Luca Manfredil, Maria Santucci de Magistrism, Maria-Jose Sánchezn, Carlota Castro-Espinq, Daniel Rodríguez Palacioss, Pilar Amianou, Marcela Guevaraw, Yvonne T. van der Schouwz, Jolanda M.A. Boeraa, W.M. Monique Verschurenz, Stephen J. Sharp, Nita G. Forouhi, Nicholas J. Wareham, Eszter P. Vamos, Kiara Chang, Paolo Vineis, Alicia K. Heath, Marc J. Gunter, Geneviève Nicolas, Elisabete Weiderpass, Inge Huybrechts, Rachel L. Batterham, Food consumption by degree of food processing and risk of type 2 diabetes mellitus: a prospective cohort analysis of the European Prospective Investigation into Cancer and Nutrition (EPIC)

The Lancet Regional Health – Europe, 2024, Volume 46, 101043. doi: 10.1016/j.lanepe.2024.101043."

Reynalda Cordova, Vivian Viallona, Emma Fontvieillea, Laia Peruchet-Noraya, Anna Jansanaa, Karl-Heinz Wagner, Cecilie Kyrø, Anne Tjønneland, Verena Katzke, Rashmita Bajracharya, Matthias B. Schulze, Giovanna Masala, Sabina Sieri, Salvatore Panico, Fulvio Ricceri, Rosario Tumino, Jolanda M.A. Boer, W.M. Monique Verschuren, Yvonne T. van der Schouw, Paula Jakszyn, Daniel Redondo-Sánchez, Pilar Amiano, José María Huertas, Marcela Guevara, Yan Bornéy, Emily Sonestedty, Konstantinos K. Tsilidis, Christopher Millett, Alicia K. Heath, Elom K. Aglago, Dagfinn Aune, Marc J. Guntera, Pietro Ferrari, Inge Huybrechts, Heinz Freisling. Consumption of ultra-processed foods and risk of multimorbidity of cancer and cardiometabolic diseases: a multinational cohort study. The Lancet Regional Health – Europe 2024, Volume 35, 100771

Rubén Fernández-Rodríguez, Bruno Bizzozero-Peroni, Valentina Díaz-Goñi, Miriam Garrido-Miguel, Gabriele Bertotti, Alberto Roldán-Ruiz, Miguel López-Moreno. Plant-based meat alternatives and cardiometabolic health: a systematic review and meta-analysis. The American Journal of Clinical Nutrition 2025. Volume 121, Issue 2, February 2025, Pages 274-283 doi: <https://doi.org/10.1016/j.ajcnut.2024.12.002>

Messina M, Duncan AM, Glenn AJ, Mariotti F. Perspective: Plant-Based Meat Alternatives Can Help Facilitate and Maintain a Lower Animal to Plant Protein Intake Ratio. Adv Nutr. 2023 May;14(3):392-405. doi: 10.1016/j.advnut.2023.03.003. Epub 2023 Mar 10. PMID: 36906147; PMCID: PMC10201680.

Page 6

Sarah Nájera Espinosa, Genevieve Hadida, Anne Jelmar Sietsma, Carmelia Alae-Carew, Grace Turner, Rosemary Green, Silvia Pastorino, Roberto Picetti, Pauline Scheelbeek, Mapping the evidence of novel plant-based foods: a systematic review of nutritional, health, and environmental impacts in high-income countries, Nutrition Reviews, 2024;, nuae031, doi: <https://doi.org/10.1093/nutrit/nuae031>

Cara L. Cargo-Froom, Rex W. Newkirk, Christopher P.F. Marinangeli, Anna K. Shoveller, Yongfeng Ai, and Daniel A. Columbus. 2023. The effects of extrusion on nutrient content of Canadian pulses with a focus on protein and amino acids. *Canadian Journal of Animal Science*. 103(1): 44-58. <https://doi.org/10.1139/cjas-2022-0088>

Samtiya, M., Aluko, R.E. & Dhewa, T. Plant food anti-nutritional factors and their reduction strategies: an overview. *Food Prod Process and Nutr* 2, 6 (2020). <https://doi.org/10.1186/s43014-020-0020-5>

Xia S, Song J, Ma C, Hao T, Hou Y, Shen S, Li Z, Xue Y, Xue C, Jiang X. A mechanistic investigation into combined influences of NaCl and extrusion temperature on fibrous structures of high-moisture textured yeast protein. *Food Chem*. 2024 Dec 1;460(Pt 1):140421. doi: 10.1016/j.foodchem.2024.140421. Epub 2024 Jul 14. PMID: 39032293.

Page 9

Ian F Pryme, Truls M Aarra. Exhaustive Overview of Dietary Plant Lectins: Prospective Importance in the Mediterranean Diet. *Am J Biomed Sci & Res*. 2021 - 13(4). AJBSR.MS.ID.001883. DOI: 10.34297/AJBSR.2021.13.001883.

Radhakrishnan, A., Chellapandian, H., Ramasamy, P. et al. Back2Basics: animal lectins: an insight into a highly versatile recognition protein. *J Proteins Proteom* 14, 43–59 (2023). <https://doi.org/10.1007/s42485-022-00102-4>

Page 10

Gréa C, Dittmann A, Wolff D, Werner R, Turban C, Roser S, Hoffmann I, Storcksdieck genannt Bonsmann S. Comparison of the Declared Nutrient Content of Plant-Based Meat Substitutes and Corresponding Meat Products and Sausages in Germany. *Nutrients*. 2023; 15(18):3864. <https://doi.org/10.3390/nu15183864>

van Haperen, Martine. 2023 “HOE GEZOND ZIJN VLEESVERVANGERS?” Accessed April 1, 2025. https://proveg.com/nl/wp-content/uploads/sites/6/2023/08/PV_NL_Vleesvervangers-Rapport_final.pdf

Sara de las Heras-Delgado, Sangeetha Shyam, Èrica Cunillera, Natalia Dragusan, Jordi Salas-Salvadó, Nancy Babio. Are plant-based alternatives healthier? A two-dimensional evaluation from nutritional and processing standpoints. *Food Research International*. Volume 169, July 2023, 112857.

Bryngelsson, Susanne, Hanieh Moshtaghian, Marta Bianchi, and Elinor Hallström. 2022. “Nutritional Assessment of Plant-Based Meat Analogues on the Swedish Market.” *International Journal of Food Sciences and Nutrition* 73 (7): 889–901.

Alessandrini, Roberta, Mhairi K. Brown, Sonia Pombo-Rodrigues, Sheena Bhageerutty, Feng J. He, and Graham A. MacGregor. 2021. “Nutritional Quality of Plant-Based Meat Products Available in the UK: A Cross-Sectional Survey.” *Nutrients* 13 (12). <https://doi.org/10.3390/nu13124225>.

Page 11

Neill, H. R., Gill, C. I. R., McDonald, E. J., McRoberts, W. C., & Pourshahidi, L. K. (2021). The future is bright: Biofortification of common foods can improve vitamin D status. *Critical Reviews in Food Science and Nutrition*, 63(4), 505–521. <https://doi.org/10.1080/10408398.2021.1950609>

ADHB. Vitamins & minerals for beef and sheep farmers. Available from: <https://ahdb.org.uk/knowledge-library/vitamins-minerals-for-beef-and-sheep-farmers> Accessed: 1 April 2025.

Pork gateway. Growing-Finishing Swine Nutrient Recommendations and Feeding Management. Available from: <https://porkgateway.org/resource/growing-finishing-swine-nutrient-recommendations-and-feeding-management/> Accessed: 1 April 2025.

FAO. Regulation on the fortification of certain foods (LIVSFS 2018:5). Available from: <https://www.fao.org/faolex/results/details/en/c/LEX-FAOC202204/> Accessed: 1 April 2025.

European Commission. Questions and Answers on Fortified Foods. Available from: https://ec.europa.eu/commission/presscorner/detail/en/memo_06_199 Accessed: 1 April 2025.

Voedingcentrum. Schijf van Vijf-vak: peulvruchten, noten, vis, ei, vlees en zuivel(alternatieven). Available from: <https://www.voedingcentrum.nl/nl/gezond-eten-met-de-schijf-van-vijf/wat-staat-in-de-schijf-van-vijf-en-wat-niet/vis-peulvruchten-vlees-ei-noten-en-zuivel.aspx> Accessed: 1 April 2025.

Valentina Gallani, Anna-Lena Klapp, Building bridges between habit and health, Proveg 2024. Available from: https://proveg.org/wp-content/uploads/2024/11/INT_Research_Plant-based-alternatives-comparison_2024-1.pdf

Page 12

Monteiro CA, Cannon G, Levy RB, Moubarac JC, Louzada ML, Rauber F, Khandpur N, Cediel G, Neri D, Martinez-Steele E, Baraldi LG, Jaime PC. Ultra-processed foods: what they are and how to identify them. *Public Health Nutr.* 2019 Apr;22(5):936-941. doi: 10.1017/S1368980018003762. Epub 2019 Feb 12. PMID: 30744710; PMCID: PMC10260459.

Page 16

Fernanda Rauber, Maria Laura da Costa Louzadab, Kiara Chang, Inge Huybrechtse, Marc J. Gunter, Carlos Augusto Monteiro, Eszter P. Vamos, Renata Bertazzi Levy. Implications of food ultra-processing on cardiovascular risk considering plant origin foods: an analysis of the UK Biobank cohort. *The Lancet Regional Health – Europe* 2024. Volume 43, 100948.

Hall et al NCT 05290064

Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, Chung ST, Costa E, Courville A, Darcey V, Fletcher LA, Forde CG, Gharib AM, Guo J, Howard R, Joseph PV, McGehee S, Ouwerkerk R, Raising K, Rozga I, Stagliano M, Walter M, Walter PJ, Yang S, Zhou M. Ultra-Processed Diets Cause Excess Calorie Intake and Weight Gain: An Inpatient Randomized Controlled Trial of Ad Libitum Food Intake. *Cell Metab.* 2019 Jul 2;30(1):67-77.e3. doi: 10.1016/j.cmet.2019.05.008. Epub 2019 May 16. Erratum in: *Cell Metab.* 2019 Jul 2;30(1):226. doi: 10.1016/j.cmet.2019.05.020. Erratum in: *Cell Metab.* 2020 Oct 6;32(4):690. doi: 10.1016/j.cmet.2020.08.014. PMID: 31105044; PMCID: PMC7946062.

Hamano S, Sawada M, Aihara M, Sakurai Y, Sekine R, Usami S, Kubota N, Yamauchi T. Ultra-processed foods cause weight gain and increased energy intake associated with reduced chewing frequency: A randomized, open-label, crossover study. *Diabetes Obes Metab.* 2024 Nov;26(11):5431-5443. doi: 10.1111/dom.15922. Epub 2024 Sep 12. PMID: 39267249.

Page 18

Mills, S., Brown, H., Wrieden, W. et al. Frequency of eating home cooked meals and potential benefits for diet and health: cross-sectional analysis of a population-based cohort study. *Int J Behav Nutr Phys Act* 14, 109 (2017). <https://doi.org/10.1186/s12966-017-0567-y>

Bajunaid, R., Niu, C., Hambly, C. et al. Predictive equation derived from 6,497 doubly labelled water measurements enables the detection of erroneous self-reported energy intake. *Nat Food* 6, 58–71 (2025). <https://doi.org/10.1038/s43016-024-01089-5>

Mavlanov U, Czaja TP, Nuriddinov S, Dalimova D, Dragsted LO, Engelsens SB, Khakimov B. The effects of industrial processing and home cooking practices on trans-fatty acid profiles of vegetable oils. *Food Chem.* 2025 Mar 30;469:142571. doi: 10.1016/j.foodchem.2024.142571. Epub 2024 Dec 18. PMID: 39721446.

Lee S, Choi Y, Jeong HS, Lee J, Sung J. Effect of different cooking methods on the content of vitamins and true retention in selected vegetables. *Food Sci Biotechnol.* 2017 Dec 12;27(2):333-342. doi: 10.1007/s10068-017-0281-1. PMID: 30263756; PMCID: PMC6049644.

Davidou S , Christodoulou A , Fardet A , Frank K . The holistico-reductionist Siga classification according to the degree of food processing: an evaluation of ultra-processed foods in French supermarkets. *Food Funct.* 2020 Mar 1;11(3):2026-2039. doi: 10.1039/c9fo02271f. Epub 2020 Feb 21. PMID: 32083627.

Ahrné, L., Chen, H., Henry, C.J. et al. Defining the role of processing in food classification systems—the IUFoST formulation & processing approach. *npj Sci Food* 9, 56 (2025). <https://doi.org/10.1038/s41538-025-00395-x>

Mills, S., Brown, H., Wrieden, W. et al. Frequency of eating home cooked meals and potential benefits for diet and health: cross-sectional analysis of a population-based cohort study. *Int J Behav Nutr Phys Act* 14, 109 (2017). <https://doi.org/10.1186/s12966-017-0567-y>

Bajunaid, R., Niu, C., Hambly, C. et al. Predictive equation derived from 6,497 doubly labelled water measurements enables the detection of erroneous self-reported energy intake. *Nat Food* 6, 58–71 (2025). <https://doi.org/10.1038/s43016-024-01089-5>

Mavlanov U, Czaja TP, Nuriddinov S, Dalimova D, Dragsted LO, Engelsen SB, Khakimov B. The effects of industrial processing and home cooking practices on trans-fatty acid profiles of vegetable oils. *Food Chem.* 2025 Mar 30;469:142571. doi: 10.1016/j.foodchem.2024.142571. Epub 2024 Dec 18. PMID: 39721446.

Lee S, Choi Y, Jeong HS, Lee J, Sung J. Effect of different cooking methods on the content of vitamins and true retention in selected vegetables. *Food Sci Biotechnol.* 2017 Dec 12;27(2):333-342. doi: 10.1007/s10068-017-0281-1. PMID: 30263756; PMCID: PMC6049644.

Page 20

Heras-Delgado S. et al. Are plant-based alternatives healthier? A two-dimensional evaluation from nutritional and processing standpoints *Food Research International.* 169:112875 (July 2023)

Sarda B, Kesse-Guyot E, Deschamps V, et al. Complementarity between the updated version of the front-of-pack nutrition label Nutri-Score and the food-processing NOVA classification. *Public Health Nutrition.* 2024;27(1):e63. doi:10.1017/S1368980024000296

Page 21

Sarah Nájera Espinosa, Genevieve Hadida, Anne Jelmar Sietsma, Carmelia Alae-Carew, Grace Turner, Rosemary Green, Silvia Pastorino, Roberto Picetti, Pauline Scheelbeek, Mapping the evidence of novel plant-based foods: a systematic review of nutritional, health, and environmental impacts in high-income countries, *Nutrition Reviews*, 2024;, nuae031, doi: <https://doi.org/10.1093/nutrit/nuae031>

Rubén Fernández-Rodríguez, Bruno Bizzozero-Peroni, Valentina Díaz-Goñi, Miriam Garrido-Miguel, Gabriele Bertotti, Alberto Roldán-Ruiz, Miguel López-Moreno. Plant-based meat alternatives and cardiometabolic health: a systematic review and meta-analysis. *The American Journal of Clinical Nutrition* 2025. Volume 121, Issue 2, February 2025, Pages 274-283 doi: <https://doi.org/10.1016/j.ajcnut.2024.12.002>

Farsi DN, Gallegos JL, Koutsidis G, Nelson A, Finnigan TJA, Cheung W, Muñoz-Muñoz JL, Commane DM. Substituting meat for mycoprotein reduces genotoxicity and increases the abundance of beneficial microbes in the gut: Mycomeat, a randomised crossover control trial. *Eur J Nutr.* 2023 Apr;62(3):1479-1492. doi: 10.1007/s00394-023-03088-x. Epub 2023 Jan 18. PMID: 36651990; PMCID: PMC10030420.

Bottin JH, Swann JR, Cropp E, Chambers ES, Ford HE, Ghatei MA, Frost GS. Mycoprotein reduces energy intake and postprandial insulin release without altering glucagon-like peptide-1 and peptide tyrosine-tyrosine concentrations in healthy overweight and obese adults: a randomised-controlled trial. *Br J Nutr.* 2016 Jul;116(2):360-74. doi: 10.1017/S0007114516001872. Epub 2016 May 20. PMID: 27198187; PMCID: PMC4910676.

Hartley L, May MD, Loveman E, Colquitt JL, Rees K. Dietary fibre for the primary prevention of cardiovascular disease. Cochrane Database of Systematic Reviews 2016, Issue 1. Art. No.: CD011472. DOI: 10.1002/14651858.CD011472.pub2. Accessed 04 April 2025.

Hooper L, Martin N, Jimoh OF, Kirk C, Foster E, Abdelhamid AS. Reduction in saturated fat intake for cardiovascular disease. Cochrane Database of Systematic Reviews 2020, Issue 5. Art. No.: CD011737. DOI: 10.1002/14651858.CD011737.pub2. Accessed 04 April 2025.

Page 22

Reuzé A, Méjean C, Sirieix L, Baudry J, Kesse-Guyot E, Druetne-Pecollo N, Brunin J, Hercberg S, Touvier M, Péneau S, Allès B. Stages of Change toward Meat Reduction: Associations with Motives and Longitudinal Dietary Data on Animal-Based and Plant-Based Food Intakes in French Adults. *J Nutr.* 2023 Nov;153(11):3295-3307. doi: 10.1016/j.tjnut.2023.09.017. Epub 2023 Sep 28. PMID: 37774842.

GFI Europe. UK plant-based food retail market insights 2022 to 2023 with initial insights into the 2024 market. 2024. Available from: <https://gfiurope.org/wp-content/uploads/2024/10/UK-plant-based-food-retail-market-insights-October-2024.pdf>. Accessed: 04 April 2025.

Alae-Carew C, Green R, Stewart C, Cook B, Dangour AD, Scheelbeek PFD. The role of plant-based alternative foods in sustainable and healthy food systems: Consumption trends in the UK. *Sci Total Environ.* 2022 Feb 10;807(Pt 3):151041. doi: 10.1016/j.scitotenv.2021.151041. Epub 2021 Oct 18. PMID: 34673070; PMCID: PMC8724617.

Rubén Fernández-Rodríguez, Bruno Bizzozero-Peroni, Valentina Díaz-Goñi, Miriam Garrido-Miguel, Gabriele Bertotti, Alberto Roldán-Ruiz, Miguel López-Moreno. Plant-based meat alternatives and cardiometabolic health: a systematic review and meta-analysis. *The American Journal of Clinical Nutrition* 2025. Volume 121, Issue 2, February 2025, Pages 274-283 doi: <https://doi.org/10.1016/j.ajcnut.2024.12.002>

Sarah Nájera Espinosa, Genevieve Hadida, Anne Jelmar Sietsma, Carmelia Alae-Carew, Grace Turner, Rosemary Green, Silvia Pastorino, Roberto Picetti, Pauline Scheelbeek, Mapping the evidence of novel plant-based foods: a systematic review of nutritional, health, and environmental impacts in high-income countries, *Nutrition Reviews*, 2024;, nuae031, doi: <https://doi.org/10.1093/nutrit/nuae031>

Nagra M, Tsam F, Ward S, Ur E. Animal vs Plant-Based Meat: A Hearty Debate. *Can J Cardiol.* 2024 Jul;40(7):1198-1209. doi: 10.1016/j.cjca.2023.11.005. Epub 2024 Jun 26. PMID: 38934982.

Page 23

Domić J, Pinckaers PJ, Grootswagers P, Siebelink E, Gerdessen JC, van Loon LJ, de Groot LC. A Well-Balanced Vegan Diet Does not Compromise Daily Mixed Muscle Protein Synthesis Rates when Compared with an Omnivorous Diet in Active Older Adults: A Randomized Controlled Cross-Over Trial. *J Nutr.* 2024 Dec 26:S0022-3166(24)01246-X. doi: 10.1016/j.tjnut.2024.12.019. Epub ahead of print. PMID: 39732437.

Pinckaers PJ, Domić J, Petrick HL, Holwerda AM, Trommelen J, Hendriks FK, Houben LH, Goessens JP, van Kranenburg JM, Senden JM, de Groot LC, Verdijk LB, Snijders T, van Loon LJ. Higher Muscle Protein Synthesis Rates Following Ingestion of an Omnivorous Meal Compared with an Isocaloric and Isonitrogenous Vegan Meal in Healthy, Older Adults. *J Nutr.* 2024 Jul;154(7):2120-2132. doi: 10.1016/j.tjnut.2023.11.004. Epub 2023 Nov 15. PMID: 37972895.

Farsi DN, Gallegos JL, Koutsidis G, Nelson A, Finnigan TJA, Cheung W, Muñoz-Muñoz JL, Commane DM. Substituting meat for mycoprotein reduces genotoxicity and increases the abundance of beneficial microbes in the gut: Mycomeat, a randomised crossover control trial. *Eur J Nutr.* 2023 Apr;62(3):1479-1492. doi: 10.1007/s00394-023-03088-x. Epub 2023 Jan 18. PMID: 36651990; PMCID: PMC10030420.

Rubén Fernández-Rodríguez, Bruno Bizzozero-Peroni, Valentina Díaz-Goñi, Miriam Garrido-Miguel, Gabriele Bertotti, Alberto Roldán-Ruiz, Miguel López-Moreno. Plant-based meat alternatives and cardiometabolic health: a systematic review and meta-analysis. *The American Journal of Clinical Nutrition* 2025. Volume 121, Issue 2, February 2025, Pages 274-283 doi: <https://doi.org/10.1016/j.ajcnut.2024.12.002>

Schoeneck M, Iggman D. The effects of foods on LDL cholesterol levels: A systematic review of the accumulated evidence from systematic reviews and meta-analyses of randomized controlled trials. *Nutr Metab Cardiovasc Dis*. 2021 May 6;31(5):1325-1338. doi: 10.1016/j.numecd.2020.12.032. Epub 2021 Jan 16. PMID: 33762150.

2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *European Heart Journal*. 2019. doi:10.1093/eurheartj/ehz455.

Penson, P.E., Pirro, M. & Banach, M. LDL-C: lower is better for longer—even at low risk. *BMC Med* 18, 320 (2020). <https://doi.org/10.1186/s12916-020-01792-7>.

Page 25

Bianchi F, Stewart C, Astbury NM, Cook B, Aveyard P, Jebb SA. Replacing meat with alternative plant-based products (RE-MAP): a randomized controlled trial of a multicomponent behavioral intervention to reduce meat consumption. *Am J Clin Nutr*. 2022 May 1;115(5):1357-1366. doi: 10.1093/ajcn/nqab414. PMID: 34958364; PMCID: PMC9071457.

Page 26

Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, Chung ST, Costa E, Courville A, Darcey V, Fletcher LA, Forde CG, Gharib AM, Guo J, Howard R, Joseph PV, McGehee S, Ouwerkerk R, Rasinger K, Rozga I, Stagliano M, Walter M, Walter PJ, Yang S, Zhou M. Ultra-Processed Diets Cause Excess Calorie Intake and Weight Gain: An Inpatient Randomized Controlled Trial of Ad Libitum Food Intake. *Cell Metab*. 2019 Jul 2;30(1):67-77.e3. doi: 10.1016/j.cmet.2019.05.008. Epub 2019 May 16. Erratum in: *Cell Metab*. 2019 Jul 2;30(1):226. doi: 10.1016/j.cmet.2019.05.020. Erratum in: *Cell Metab*. 2020 Oct 6;32(4):690. doi: 10.1016/j.cmet.2020.08.014. PMID: 31105044; PMCID: PMC7946062.

Hamano S, Sawada M, Aihara M, Sakurai Y, Sekine R, Usami S, Kubota N, Yamauchi T. Ultra-processed foods cause weight gain and increased energy intake associated with reduced chewing frequency: A randomized, open-label, crossover study. *Diabetes Obes Metab*. 2024 Nov;26(11):5431-5443. doi: 10.1111/dom.15922. Epub 2024 Sep 12. PMID: 39267249.

Page 27

Fernanda Rauber, Maria Laura da Costa Louzadab, Kiara Chang, Inge Huybrechtse, Marc J. Gunter, Carlos Augusto Monteiro, Eszter P. Vamos, Renata Bertazzi Levy. Implications of food ultra-processing on cardiovascular risk considering plant origin foods: an analysis of the UK Biobank cohort. *The Lancet Regional Health – Europe* 2024. Volume 43, 100948.

Chang K, Parnham JC, Rauber F, Levy RB, Huybrechts I, Gunter MJ, Millett C, Vamos EP. Plant-based dietary patterns and ultra-processed food consumption: a cross-sectional analysis of the UK Biobank. *EClinicalMedicine*. 2024 Nov 14;78:102931. doi: 10.1016/j.eclinm.2024.102931. PMID: 39619237; PMCID: PMC11605131.

Parra-Soto S, Ahumada D, Petermann-Rocha F, Boonpoor J, Gallegos JL, Anderson J, Sharp L, Malcomson FC, Livingstone KM, Mathers JC, Pell JP, Ho FK, Celis-Morales C. Association of meat, vegetarian, pescatarian and fish-poultry diets with risk of 19 cancer sites and all cancer: findings from the UK Biobank prospective cohort study and meta-analysis. *BMC Med*. 2022 Feb 24;20(1):79. doi: 10.1186/s12916-022-02257-9. PMID: 35655214; PMCID: PMC9281076.

Petermann-Rocha F, Parra-Soto S, Gray S, Anderson J, Welsh P, Gill J, Sattar N, Ho FK, Celis-Morales C, Pell JP. Vegetarians, fish, poultry, and meat-eaters: who has higher risk of cardiovascular disease incidence and mortality? A prospective study from UK Biobank. *Eur Heart J*. 2021 Mar 21;42(12):1136-1143. doi: 10.1093/eurheartj/ehaa939. PMID: 33313747; PMCID: PMC7982287.

International Agency for Research on Cancer. Epic-Europe Study. Available from: <https://epic.iarc.fr/>. Accessed: 14 April 2025.

María Seguí-Gómez, Carmen de la Fuente, Zenaida Vázquez, Jokin de Irala, Miguel A Martínez-González, Cohort profile: The ‘Seguimiento Universidad de Navarra’ (SUN) study, *International Journal of Epidemiology*, Volume 35, Issue 6, December 2006, Pages 1417–1422, <https://doi.org/10.1093/ije/dyl223>

UK Biobank. Available From: <https://www.ukbiobank.ac.uk/> Accessed:14 April 2025.

CDC. About NAHNES. Available from: <https://www.cdc.gov/nchs/nhanes/about/index.html> Accessed:14 April 2025.

Nurses' Health Study. Available from: <https://nurseshealthstudy.org/> Accessed:14 April 2025.

Page 29

Riboli E, Hunt K, Slimani N, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutrition*. 2002;5(6b):1113-1124. doi:10.1079/PHN2002394

UK Biobank. 24-hour dietary recall questionnaire (Oxford WebQ) Version 1.4. Available from:<https://biobank.ndph.ox.ac.uk/ukb/ukb/docs/DietWebQ.pdf> Accessed:14 April 2025.

"Samuel J. Dicken, Christina C. Dahm, Daniel B. Ibsen, Anja Olsen, Anne Tjønnelande, Mariem Louati-Hajjig, Claire Cadeaug, Chloé Marquesg, Matthias B. Schulzeh, Franziska Jannaschh, Ivan Baldassarik, Luca Manfredil, Maria Santucci de Magistrism, Maria-Jose Sánchezn, Carlota Castro-Espinq, Daniel Rodríguez Palacioss, Pilar Amianou, Marcela Guevaraw, Yvonne T. van der Schouwz, Jolanda M.A. Boeraa, W.M. Monique Verschurenz, Stephen J. Sharp, Nita G. Forouhi, Nicholas J. Wareham, Eszter P. Vamos, Kiara Chang, Paolo Vineis, Alicia K. Heath, Marc J. Gunter, Geneviève Nicolas, Elisabete Weiderpass, Inge Huybrechts, Rachel L. Batterham, Food consumption by degree of food processing and risk of type 2 diabetes mellitus: a prospective cohort analysis of the European Prospective Investigation into Cancer and Nutrition (EPIC)

The Lancet Regional Health – Europe, 2024, Volume 46, 101043. doi: 10.1016/j.lanepe.2024.101043."

Chang K, Gunter MJ, Rauber F, Levy RB, Huybrechts I, Kliemann N, Millett C, Vamos EP. Ultra-processed food consumption, cancer risk and cancer mortality: a large-scale prospective analysis within the UK Biobank. *EClinicalMedicine*. 2023 Jan 31;56:101840. doi: 10.1016/j.eclinm.2023.101840. PMID: 36880051; PMCID: PMC9985039.

Nguyen M, Jarvis SE, Tinajero MG, Yu J, Chiavaroli L, Mejia SB, Khan TA, Tobias DK, Willett WC, Hu FB, Hanley AJ, Birken CS, Sievenpiper JL, Malik VS. Sugar-sweetened beverage consumption and weight gain in children and adults: a systematic review and meta-analysis of prospective cohort studies and randomized controlled trials. *Am J Clin Nutr*. 2023 Jan;117(1):160-174. doi: 10.1016/j.ajcnut.2022.11.008. Epub 2022 Dec 20. PMID: 36789935.

Feng Q, Wong SH, Zheng J, Yang Q, Sung JJ, Tsoi KK. Intake of processed meat, but not sodium, is associated with risk of colorectal cancer: Evidence from a large prospective cohort and two-sample Mendelian randomization. *Clin Nutr*. 2021 Jul;40(7):4551-4559. doi: 10.1016/j.clnu.2021.05.036. Epub 2021 Jun 14. PMID: 34229259.

Page 30

Mendoza K, Smith-Warner SA, Rossato SL, Khandpur N, Manson JE, Qi L, Rimm EB, Mukamal KJ, Willett WC, Wang M, Hu FB, Mattei J, Sun Q. Ultra-processed foods and cardiovascular disease: analysis of three large US prospective cohorts and a systematic review and meta-analysis of prospective cohort studies. *Lancet Reg Health Am*. 2024 Sep 2;37:100859. doi: 10.1016/j.lana.2024.100859. PMID: 39286398; PMCID: PMC11403639.

Cordova R, Viallon V, Fontvieille E, Peruchet-Noray L, Jansana A, Wagner KH, Kyrø C, Tjønneland A, Katzke V, Bajracharya R, Schulze MB, Masala G, Sieri S, Panico S, Ricceri F, Tumino R, Boer JMA, Verschuren WMM, van der Schouw YT, Jakšzyn P, Redondo-Sánchez D, Amiano P, Huerta JM, Guevara M, Borné Y, Sonestedt E, Tsilidis KK, Millett C, Heath AK, Aglago EK, Aune D, Gunter MJ, Ferrari P, Huybrechts I, Freisling H. Consumption of ultra-processed foods and risk of multimorbidity of cancer and cardiometabolic diseases: a multinational cohort study. *Lancet Reg Health Eur*. 2023 Nov 14;35:100771. doi: 10.1016/j.lanepe.2023.100771. PMID: 38115963; PMCID: PMC10730313.

Samuel J. Dicken, Christina C. Dahm, Daniel B. Ibsen, Anja Olsen, Anne Tjønnelande, Mariem Louati-Hajjig, Claire Cadeaug, Chloé Marquesg, Matthias B. Schulzeh, Franziska Jannaschh, Ivan Baldassarik, Luca Manfredil, Maria Santucci de Magistrism, Maria-Jose Sánchezn, Carlota Castro-Espinq, Daniel Rodríguez Palacioss, Pilar Amianou, Marcela Guevaraw, Yvonne T. van der Schouwz, Jolanda M.A. Boeraa, W.M. Monique Verschurenz, Stephen J. Sharp, Nita G. Forouhi, Nicholas J. Wareham, Eszter P. Vamos, Kiara Chang, Paolo Vineis, Alicia K. Heath, Marc J. Gunter, Geneviève Nicolas, Elisabete Weiderpass, Inge Huybrechts, Rachel L. Batterham, Food consumption by degree of food processing and risk of type 2 diabetes mellitus: a prospective cohort analysis of the European Prospective Investigation into Cancer and Nutrition (EPIC)

The Lancet Regional Health – Europe, 2024, Volume 46, 101043. doi: 10.1016/j.lanepe.2024.101043."

Mendoza K, Barquera S, Tobias DK. Targeting ultra-processed foods for prevention of type 2 diabetes: state of the evidence and future directions. *Diabetologia*. 2025 Mar;68(3):495-506. doi: 10.1007/s00125-025-06358-x. Epub 2025 Jan 18. PMID: 39825911.

Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, Chung ST, Costa E, Courville A, Darcey V, Fletcher LA, Forde CG, Gharib AM, Guo J, Howard R, Joseph PV, McGehee S, Ouwerkerk R, Raisinger K, Rozga I, Stagliano M, Walter M, Walter PJ, Yang S, Zhou M. Ultra-Processed Diets Cause Excess Calorie Intake and Weight Gain: An Inpatient Randomized Controlled Trial of Ad Libitum Food Intake. *Cell Metab*. 2019 Jul 2;30(1):67-77.e3. doi: 10.1016/j.cmet.2019.05.008. Epub 2019 May 16. Erratum in: *Cell Metab*. 2019 Jul 2;30(1):226. doi: 10.1016/j.cmet.2019.05.020. Erratum in: *Cell Metab*. 2020 Oct 6;32(4):690. doi: 10.1016/j.cmet.2020.08.014. PMID: 31105044; PMCID: PMC7946062.

Hamano S, Sawada M, Aihara M, Sakurai Y, Sekine R, Usami S, Kubota N, Yamauchi T. Ultra-processed foods cause weight gain and increased energy intake associated with reduced chewing frequency: A randomized, open-label, crossover study. *Diabetes Obes Metab*. 2024 Nov;26(11):5431-5443. doi: 10.1111/dom.15922. Epub 2024 Sep 12. PMID: 39267249.

Page 32

Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, Chung ST, Costa E, Courville A, Darcey V, Fletcher LA, Forde CG, Gharib AM, Guo J, Howard R, Joseph PV, McGehee S, Ouwerkerk R, Raisinger K, Rozga I, Stagliano M, Walter M, Walter PJ, Yang S, Zhou M. Ultra-Processed Diets Cause Excess Calorie Intake and Weight Gain: An Inpatient Randomized Controlled Trial of Ad Libitum Food Intake. *Cell Metab*. 2019 Jul 2;30(1):67-77.e3. doi: 10.1016/j.cmet.2019.05.008. Epub 2019 May 16. Erratum in: *Cell Metab*. 2019 Jul 2;30(1):226. doi: 10.1016/j.cmet.2019.05.020. Erratum in: *Cell Metab*. 2020 Oct 6;32(4):690. doi: 10.1016/j.cmet.2020.08.014. PMID: 31105044; PMCID: PMC7946062.

Hall KD, Guo J, Courville AB, Boring J, Brychta R, Chen KY, Darcey V, Forde CG, Gharib AM, Gallagher I, Howard R, Joseph PV, Milley L, Ouwerkerk R, Raisinger K, Rozga I, Schick A, Stagliano M, Torres S, Walter M, Walter P, Yang S, Chung ST. Effect of a plant-based, low-fat diet versus an animal-based, ketogenic diet on ad libitum energy intake. *Nat Med*. 2021 Feb;27(2):344-353. doi: 10.1038/s41591-020-01209-1. Epub 2021 Jan 21. PMID: 33479499.

Page 33

Hamano S, Sawada M, Aihara M, Sakurai Y, Sekine R, Usami S, Kubota N, Yamauchi T. Ultra-processed foods cause weight gain and increased energy intake associated with reduced chewing frequency: A randomized, open-label, crossover study. *Diabetes Obes Metab*. 2024 Nov;26(11):5431-5443. doi: 10.1111/dom.15922. Epub 2024 Sep 12. PMID: 39267249.

ANSES. Aliments dits ultratransformés : mieux comprendre leurs effets potentiels sur la santé (2025). Available from: <https://www.anses.fr/fr/content/aliments-dits-ultratransformes-mieux-comprendre-leurs-effets-potentiels-sur-la-sante>. Accessed 14 April 2025.

Bröder J, Tauer J, Liaskos M, Hieronimus B für die Arbeitsgruppe „(Stark) verarbeitete Lebensmittel“ der DGE: Verzehr stark verarbeiteter Lebensmittel und ernährungsmitbedingte Erkrankungen: Eine systematische

Übersichtsarbeit. In: Deutsche Gesellschaft für Ernährung (Hrsg.): 15. DGE-Ernährungsbericht. Vorveröffentlichung Kapitel 9. Bonn (2023) V2–V40 <https://www.dge.de/fileadmin/dok/wissenschaft/ernaehrungsberichte/15eb/15-DGE-EB-Vorveroeffentlichung-Kapitel9.pdf>

SACN. SACN statement on processed foods and health (2023) and Available from: <https://assets.publishing.service.gov.uk/media/64ac1fe7b504f7000ccdb89a/SACN-position-statement-Processed-Foods-and-Health.pdf> Accessed 14 April 2025.

SACN. Processed foods and health: SACN's rapid evidence update summary (2025) Available from: <https://www.gov.uk/government/publications/processed-foods-and-health-sacns-rapid-evidence-update/processed-foods-and-health-sacns-rapid-evidence-update-summary> Accessed 14 April 2025.

Food, Diet and Obesity Committee. Recipe for health: a plan to fix our broken food system (2024). Available from: <https://publications.parliament.uk/pa/ld5901/ldselect/ldmfdo/19/1902.htm>. Accessed 14 April 2025.

AESAN. Revista del Comité Científico de la AESAN No. 31 (2020). Available from: https://www.aesan.gob.es/AECOSAN/docs/documentos/publicaciones/revistas_comite_cientifico/comite_cientifico_31.pdf Accessed 14 April 2025.

Nordic Council of Ministers. Nordic Nutrition Recommendations 2023. Available from: <https://www.norden.org/en/publication/nordic-nutrition-recommendations-2023> Accessed 14 April 2025.

Page 36

Feng Q, Wong SH, Zheng J, Yang Q, Sung JJ, Tsoi KK. Intake of processed meat, but not sodium, is associated with risk of colorectal cancer: Evidence from a large prospective cohort and two-sample Mendelian randomization. Clin Nutr. 2021 Jul;40(7):4551-4559. doi: 10.1016/j.clnu.2021.05.036. Epub 2021 Jun 14. PMID: 34229259.

Kelbrick M, Hesse E, O' Brien S. Cultivating antimicrobial resistance: how intensive agriculture ploughs the way for antibiotic resistance. Microbiology (Reading). 2023 Aug;169(8):001384. doi: 10.1099/mic.0.001384. PMID: 37606636; PMCID: PMC10482381.

Hayek MN. The infectious disease trap of animal agriculture. Sci Adv. 2022 Nov 4;8(44):eadd6681. doi: 10.1126/sciadv.add6681. Epub 2022 Nov 2. PMID: 36322670; PMCID: PMC9629715.

Page 41

Quorn, The production of alternative first-class protein source for a balanced, sustainable diet. White paper (2017). Available from: <https://docslib.org/download/10116636/quorn-the-production-of-alternative-first-class-protein-source-for-a-balanced-sustainable-diet>. Accessed 29 April 2025.

Robinson E, Khuttan M, McFarland-Lesser I, Patel Z, Jones A. Calorie reformulation: a systematic review and meta-analysis examining the effect of manipulating food energy density on daily energy intake. *Int J Behav Nutr Phys Act*. 2022 Apr 22;19(1):48. doi: 10.1186/s12966-022-01287-z. PMID: 35459185; PMCID: PMC9026919.

Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, Chung ST, Costa E, Courville A, Darcey V, Fletcher LA, Forde CG, Gharib AM, Guo J, Howard R, Joseph PV, McGehee S, Ouwerkerk R, Raisinger K, Rozga I, Stagliano M, Walter M, Walter PJ, Yang S, Zhou M. Ultra-Processed Diets Cause Excess Calorie Intake and Weight Gain: An Inpatient Randomized Controlled Trial of Ad Libitum Food Intake. *Cell Metab*. 2019 Jul 2;30(1):67-77.e3. doi: 10.1016/j.cmet.2019.05.008. Epub 2019 May 16. Erratum in: *Cell Metab*. 2019 Jul 2;30(1):226. doi: 10.1016/j.cmet.2019.05.020. Erratum in: *Cell Metab*. 2020 Oct 6;32(4):690. doi: 10.1016/j.cmet.2020.08.014. PMID: 31105044; PMCID: PMC7946062.

Hamano S, Sawada M, Aihara M, Sakurai Y, Sekine R, Usami S, Kubota N, Yamauchi T. Ultra-processed foods cause weight gain and increased energy intake associated with reduced chewing frequency: A randomized, open-label, crossover study. *Diabetes Obes Metab*. 2024 Nov;26(11):5431-5443. doi: 10.1111/dom.15922. Epub 2024 Sep 12. PMID: 39267249.

Gréa C, Dittmann A, Wolff D, Werner R, Turban C, Roser S, Hoffmann I, Storcksdieck genannt Bonsmann S. Comparison of the Declared Nutrient Content of Plant-Based Meat Substitutes and Corresponding Meat Products and Sausages in Germany. *Nutrients*. 2023; 15(18):3864. <https://doi.org/10.3390/nu15183864>

van Haperen, Martine. 2023 “HOE GEZOND ZIJN VLEESVERVANGERS?” Available from: https://proveg.com/nl/wp-content/uploads/sites/6/2023/08/PV_NL_Vleesvervangers-Rapport_final.pdf. Accessed April 1, 2025.

Sara de las Heras-Delgado, Sangeetha Shyam, Èrica Cunillera, Natalia Dragusan, Jordi Salas-Salvadó, Nancy Babio. Are plant-based alternatives healthier? A two-dimensional evaluation from nutritional and processing standpoints. *Food Research International*. Volume 169, July 2023, 112857.

Bryngelsson, Susanne, Hanieh Moshtaghian, Marta Bianchi, and Elinor Hallström. 2022. “Nutritional Assessment of Plant-Based Meat Analogues on the Swedish Market.” *International Journal of Food Sciences and Nutrition* 73 (7): 889–901.

Alessandrini, Roberta, Mhairi K. Brown, Sonia Pombo-Rodrigues, Sheena Bhageerutty, Feng J. He, and Graham A. MacGregor. 2021. “Nutritional Quality of Plant-Based Meat Products Available in the UK: A Cross-Sectional Survey.” *Nutrients* 13 (12). <https://doi.org/10.3390/nu13124225>.

Nesta, Targeting the health of the nation. White paper (2024). Available from: <https://www.nesta.org.uk/report/targeting-the-health-of-a-nation/identifying-the-best-measure-for-health/>. Accessed 29 April 2025.

Thompson SV, Hannon BA, An R, Holscher HD. Effects of isolated soluble fiber supplementation on body weight, glycemia, and insulinemia in adults with overweight and obesity: a systematic review and meta-analysis of randomized controlled trials. *Am J Clin Nutr*. 2017 Dec;106(6):1514-1528. doi: 10.3945/ajcn.117.163246. Epub 2017 Nov 1. PMID: 29092878.

Hartley L, May MD, Loveman E, Colquitt JL, Rees K. Dietary fibre for the primary prevention of cardiovascular disease. *Cochrane Database of Systematic Reviews* 2016, Issue 1. Art. No.: CD011472. DOI: 10.1002/14651858.CD011472.pub2

Threapleton DE, Greenwood DC, Evans CE, Cleghorn CL, Nykjaer C, Woodhead C, Cade JE, Gale CP, Burley VJ. Dietary fibre intake and risk of cardiovascular disease: systematic review and meta-analysis. *BMJ*. 2013 Dec 19;347:f6879. doi: 10.1136/bmj.f6879. PMID: 24355537; PMCID: PMC3898422.

Stephen AM, Champ MM-J, Cloran SJ, et al. Dietary fibre in Europe: current state of knowledge on definitions, sources, recommendations, intakes and relationships to health. *Nutrition Research Reviews*. 2017;30(2):149-190. doi:10.1017/S095442241700004X

So D, Whelan K, Rossi M, Morrison M, Holtmann G, Kelly JT, Shanahan ER, Staudacher HM, Campbell KL. Dietary fiber intervention on gut microbiota composition in healthy adults: a systematic review and meta-analysis. *Am J Clin Nutr*. 2018 Jun 1;107(6):965-983. doi: 10.1093/ajcn/nqy041. PMID: 29757343.

Manzanilla-Valdez ML, Ma Z, Mondor M, Hernández-Álvarez AJ. Decoding the Duality of Antinutrients: Assessing the Impact of Protein Extraction Methods on Plant-Based Protein Sources. *J Agric Food Chem*. 2024 Jun 5;72(22):12319-12339. doi: 10.1021/acs.jafc.4c00380. Epub 2024 May 23. PMID: 38780067; PMCID: PMC11157537.

Domić J, Pinckaers PJ, Grootswagers P, Siebelink E, Gerdessen JC, van Loon LJ, de Groot LC. A Well-Balanced Vegan Diet Does not Compromise Daily Mixed Muscle Protein Synthesis Rates when Compared with an Omnivorous Diet in Active Older Adults: A Randomized Controlled Cross-Over Trial. *J Nutr*. 2025 Apr;155(4):1141-1150. doi: 10.1016/j.tjnut.2024.12.019. Epub 2024 Dec 26. PMID: 39732437.

Jäger, R., Kerkick, C.M., Campbell, B.I. et al. International Society of Sports Nutrition Position Stand: protein and exercise. *J Int Soc Sports Nutr* 14, 20 (2017). <https://doi.org/10.1186/s12970-017-0177-8>.

Janssen TAH, Van Every DW, Phillips SM. The impact and utility of very low-calorie diets: the role of exercise and protein in preserving skeletal muscle mass. *Curr Opin Clin Nutr Metab Care*. 2023 Nov 1;26(6):521-527. doi: 10.1097/MCO.0000000000000980. Epub 2023 Sep 7. PMID: 37724991; PMCID: PMC10552824.

Page 46

Hooper L, Martin N, Jimoh OF, Kirk C, Foster E, Abdelhamid AS. Reduction in saturated fat intake for cardiovascular disease. *Cochrane Database of Systematic Reviews* 2020, Issue 8. Art. No.: CD011737. DOI: 10.1002/14651858.CD011737.pub3. Accessed 29 April 2025.

Page 47

Santos LP, Gigante DP, Delpino FM, Maciel AP, Bielemann RM. Sugar sweetened beverages intake and risk of obesity and cardiometabolic diseases in longitudinal studies: A systematic review and meta-analysis with 1.5 million individuals. *Clin Nutr ESPEN*. 2022 Oct;51:128-142. doi: 10.1016/j.clnesp.2022.08.021. Epub 2022 Aug 24. PMID: 36184197.

Huang Y, Chen Z, Chen B, Li J, Yuan X, Li J, Wang W, Dai T, Chen H, Wang Y, Wang R, Wang P, Guo J, Dong Q, Liu C, Wei Q, Cao D, Liu L. Dietary sugar consumption and health: umbrella review. *BMJ*. 2023 Apr 5;381:e071609. doi: 10.1136/bmj-2022-071609. PMID: 37019448; PMCID: PMC10074550.

Page 48

Santos LP, Gigante DP, Delpino FM, Maciel AP, Bielemann RM. Sugar sweetened beverages intake and risk of obesity and cardiometabolic diseases in longitudinal studies: A systematic review and meta-analysis with 1.5 million individuals. *Clin Nutr ESPEN*. 2022 Oct;51:128-142. doi: 10.1016/j.clnesp.2022.08.021. Epub 2022 Aug 24. PMID: 36184197.

European Commission, Defining dietary salt and sodium - dietary recommendations for salt/sodium intake in adults as described by relevant food and health-related organisations (2021). Available from: https://knowledge4policy.ec.europa.eu/health-promotion-knowledge-gateway/defining-dietary-salt-sodium-table-3a_en . Accessed 29 April 2025.

Graudal NA, Hubeck-Graudal T, Jurgens G. Effects of low sodium diet versus high sodium diet on blood pressure, renin, aldosterone, catecholamines, cholesterol, and triglyceride. *Cochrane Database of Systematic Reviews* 2020, Issue 12. Art. No.: CD004022. DOI: 10.1002/14651858.CD004022.pub5. Accessed 29 April 2025.

Rubén Fernández-Rodríguez, Bruno Bizzozero-Peroni, Valentina Díaz-Goñi, Miriam Garrido-Miguel, Gabriele Bertotti, Alberto Roldán-Ruiz, Miguel López-Moreno. Plant-based meat alternatives and cardiometabolic health: a systematic review and meta-analysis. *The American Journal of Clinical Nutrition* 2025. Volume 121, Issue 2, February 2025, Pages 274-283

Sarah Nájera Espinosa, Genevieve Hadida, Anne Jelmar Sietsma, Carmelia Alae-Carew, Grace Turner, Rosemary Green, Silvia Pastorino, Roberto Picetti, Pauline Scheelbeek, Mapping the evidence of novel plant-based foods: a systematic review of nutritional, health, and environmental impacts in high-income countries, *Nutrition Reviews*, 2024;, nuae031, doi: <https://doi.org/10.1093/nutrit/nuae031>

Sarah Nájera Espinosa, Genevieve Hadida, Anne Jelmar Sietsma, Carmelia Alae-Carew, Grace Turner, Rosemary Green, Silvia Pastorino, Roberto Picetti, Pauline Scheelbeek, Mapping the evidence of novel plant-based foods: a systematic review of nutritional, health, and environmental impacts in high-income countries, *Nutrition Reviews*, 2024;, nuae031, doi: <https://doi.org/10.1093/nutrit/nuae031>

Gréa C, Dittmann A, Wolff D, Werner R, Turban C, Roser S, Hoffmann I, Storcksdieck genannt Bonsmann S. Comparison of the Declared Nutrient Content of Plant-Based Meat Substitutes and Corresponding Meat Products and Sausages in Germany. *Nutrients*. 2023; 15(18):3864. <https://doi.org/10.3390/nu15183864>